



DRAFT REVISION OF ANNEX TO ISPM 27: DP02 - PLUM POX VIRUS (2016-007)

Summary comments

Name	Summary
Cameroon	Exame achevé
Cuba	No hay sugerencias al Protocolo de Diagnóstico de Plum Pox Virus, estamos de acuerdo con el mismo.
EPPO Σ	Finalised by the EPPO Secretariat on behalf of its 51 Member Countries.
European Union	Finalised by the European Commission on behalf of the EU and its 28 Member States on 29/09/2017.
Samoa	no further comments
South Africa	No comments from the National Plant Protection Organisation of South Africa.
Turkey	Plum pox virus -TR

#	Para	Text	Comment	SC's response
1	G	(General Comment)	Cameroon Les préoccupations que nous avons sont celles relatives à l'infrastructure et le niveau technique requis pour conduire de tels tests. Les formations et le développement de kits de diagnostic rapide pourraient aider à combler ces lacune pour les pays de notre région en général. <i>Category : TECHNICAL</i>	Noted. NPPOs are encouraged to seek suitable training from labs/experts identified in Section 6. Contact Points. However, the comment is more an implementation issue and it is outside of the TPDP's remit. It will be forwarded to the relevant IPPC bodies.
2	G	(General Comment)	Costa Rica We do not have comments on this document, we agree <i>Category : TECHNICAL</i>	NOTED
3	G	(General Comment)	Myanmar This disease absent in Myanmar. Myanmar has no comment,	NOTED

#	Para	Text	Comment	SC's response
			<i>Category : SUBSTANTIVE</i>	
4	G	(General Comment)	Peru We agree with the Draft revision of DP 2: Plum pox virus (2016-007) <i>Category : TECHNICAL</i>	NOTED
5	G	(General Comment)	United States of America Perhaps it would be more valuable if the protocol had input from researchers from a variety of regions to present more of a global expertise. <i>Category : SUBSTANTIVE</i>	Noted. The TPD discussed this draft DP revision prior to submitting it to the SC for approval for consultation.
6	G	(General Comment)	Canada Canada supports the draft diagnostic protocol on Plum pox virus (2016-007). <i>Category : SUBSTANTIVE</i>	NOTED
7	G	(General Comment)	European Union Finally, it is suggested to improve the structure of the protocol by presenting more clearly which step of the diagnostic process is involved, e.g. sampling (most likely similar for all ELISA tests and all molecular tests, respectively); sample preparation; detection of PPV (all strains); identification of PPV (universal) and identification of individual strains (including overview table as suggested before). <i>Category : TECHNICAL</i>	Modified (The DP was modified with sections identified for Host range, Symptoms, Sampling etc. Preparation of any comprehensive table may require the re-constitution of a new drafting group)
8	G	(General Comment)	Panama Panama has no comments on this document. <i>Category : EDITORIAL</i>	NOTED
9	G	(General Comment)	Saint Vincent and The Grenadines No additional comments. This standard is highly technical and would be difficult to be implemented by St. Vincent and the Grenadines <i>Category : SUBSTANTIVE</i>	NOTED NPPOs are encouraged to seek suitable training from labs/experts identified in Section 6. Contact Points. However, the comment is more an implementation issue and it is outside of the TPD's remit. It will be forwarded to the relevant IPPC bodies.
10	G	(General Comment)	Barbados Barbados has no comments to make on this document. <i>Category : SUBSTANTIVE</i>	NOTED
11	G	(General Comment)	Viet Nam Vietnam would like to request providing the	Modified

#	Para	Text	Comment	SC's response
			protocol for detection and identification of Plum pox virus using RT-LAMP (Reverse transcription - Loop-mediated isothermal amplification) method and providing the method for preserving the suspectedly infested-PPV samples/ plant sap (after extraction). Because LAMP/ RT-LAMP is rapid, accurate and cost effective for detection and identification of virus that is highly potential to apply in Vietnam, especially in Plant Quarantine Stations or In international trade, it's necessary to preserve the samples/plant sap (after extraction) after performing a diagnosis for further technical argument/discussion. <i>Category : SUBSTANTIVE</i>	A paragraph was added in Section 3. Detection and Identification that indicates that this DP describes well established methods and that other new methods such as LAMP and NGS exist, but as yet are not fully validated. A NPPO may use any suitably validated test. Preservation of plant material is out of the scope. However, some guidance for storage is provided in the Records Section of the protocol.
12	G	(General Comment)	Tajikistan I support the document as it is and I have no comments <i>Category : SUBSTANTIVE</i>	NOTED
13	G	(General Comment)	New Zealand Have no comments on the draft. <i>Category : SUBSTANTIVE</i>	NOTED
14	G	(General Comment)	Bahamas There is a need for strict plant quarantine and procedures for testing of imported nursery stock to prevent the introduction of PPV to the region. The Bahamas therefore support this diagnostic protocol. <i>Category : SUBSTANTIVE</i>	NOTED
15	G	(General Comment)	Thailand agree with the proposed draft DP for Plum pox virus <i>Category : SUBSTANTIVE</i>	NOTED
16	G	(General Comment)	Lao People's Democratic Republic Lao PDR agreed with this draft revision. <i>Category : SUBSTANTIVE</i>	NOTED
17	G	(General Comment)	Honduras HONDURAS NO TIENE COMENTARIOS <i>Category : TECHNICAL</i>	NOTED
18	G	(General Comment)	Algeria No comment <i>Category : TECHNICAL</i>	NOTED
19	G	(General Comment)	Nicaragua Nicaragua considera que este protocolo es una herramienta de gran importancia para los países, que por el comportamiento de la	NOTED

#	Para	Text	Comment	SC's response
			enfermedad tienen afectaciones del virus. El estudio de las diferentes cepas del virus permitirá contar la información actualizada para emitir un diagnóstico acertado y de manera específica sobre las diferentes cepas del virus. <i>Category : EDITORIAL</i>	
20	G	(General Comment)	Colombia No obstante, teniendo en cuenta las características variables de los síntomas que causa Plum Pox Virus –PPV-, se solicita mantener e incluir fotografías de los síntomas más relevantes, las cuales serían de gran utilidad para países que no registran esta plaga. <i>Category : TECHNICAL</i>	Modified Database links are now included for symptoms of the virus
21	33	Main discussion points during development of the diagnostic protocol	EPPO <i>Category : TECHNICAL</i>	NOTED
22	52	Sharka (plum pox) <i>Plum pox virus</i> is one of the most serious diseases of stone fruit. The disease, caused by <i>Plum pox virus</i> (PPV), affects plants of the genus <i>Prunus</i> . It is particularly detrimental in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> because it reduces quality and causes premature fruit drop. It is estimated that the costs of managing sharka worldwide since the 1970s exceed 10 000 million euros (Cambra <i>et al.</i> , 2006b).	Viet Nam Do not use an other name <i>Category : TECHNICAL</i>	Considered but not incorporated. (Sharka/Plum pox is the name of the disease. Plum pox virus is the causal agent of the disease. Under section 2 "taxonomic information" this is better explained.)
23	52	Sharka (plum pox) is one of the most serious viral diseases of stone fruit. The disease, caused by <i>Plum pox virus</i> (PPV), affects plants of the genus <i>Prunus</i> <i>Prunus</i> (family <i>Rosacea</i>). <i>Sharka was first reported in P. domestica in Bulgaria in 1917–1918, and was described as a viral disease in 1932. Since then, the virus has spread progressively to a large part of Europe, around the Mediterranean basin and the Near East. It has been found with a restricted distribution in South and North America and Asia (EPPO, 2006; CABI, 2016).</i> It is particularly detrimental in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> because it reduces quality and causes premature fruit drop. It is estimated that the costs of managing sharka worldwide since the 1970s exceed 10 000 million euros (Cambra <i>et al.</i> , 2006b).	Kenya <i>Category : SUBSTANTIVE</i>	Modified (Changes made as suggested, but some grammar and spelling corrected also)
24	52	Sharka (plum pox) is one of the most serious diseases of stone fruit. The disease, caused by <i>Plum pox virus</i> (PPV), affects plants of the genus	PPPO replace premature with early <i>Category : SUBSTANTIVE</i>	Incorporated

#	Para	Text	Comment	SC's response
		<i>Prunus</i> . It is particularly detrimental in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> because it reduces quality and causes premature <u>early</u> fruit drop. It is estimated that the costs of managing sharka worldwide since the 1970s exceed 10 000 million euros (Cambra <i>et al.</i> , 2006b).		
25	53	Sharka <i>Plum pox virus</i> was first reported in <i>P. domestica</i> in Bulgaria in 1917–1918, and was described as a viral disease in 1932. Since then, the virus has spread progressively to a large part of Europe, around the Mediterranean basin and the Near East. It has been found with a restricted distribution in South and North America and Asia (EPPO, 2006; CABI, 2016).	Viet Nam Should be use name of pest as title Category : <i>TECHNICAL</i>	Modified (Section 1 was modified. NB: Sharka/plum pox disease was first observed in 1917-1918. The viral nature was described in 1932.)
26	53	The disease Sharka was first reported is particularly detrimental in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> because it reduces quality and causes premature fruit drop. It is estimated that the costs of managing sharka worldwide since the 1970s exceed 10 000 million euros (Cambra <i>et al.</i> , 2006b). <i>P. domestica</i> in Bulgaria in 1917–1918, and was described as a viral disease in 1932. Since then, the virus has spread progressively to a large part of Europe, around the Mediterranean basin and the Near East. It has been found with a restricted distribution in South and North America and Asia (EPPO, 2006; CABI, 2016).	Kenya Category : <i>SUBSTANTIVE</i>	Modified (Section 1 was modified. NB: Sharka/plum pox disease was first observed in 1917-1918. The viral nature was described in 1932.)
27	54	<i>Plum pox virus</i> is a member of the genus <i>Potyvirus</i> in the family <i>Potyviridae</i> . The virus particles are flexuous rods of approximately 700 nm × 11 nm, and are composed of a single-stranded RNA molecule consisting of almost 10 000 nucleotides coated by up to 2 000 subunits of a single coat protein (García and Cambra, 2007). PPV is transmitted in the field by aphids in a non-persistent manner, but movement of infected propagative plant material is the main way in which PPV is spread over long distances. <u>Transmission via seed and pollen have not been reported (Pasquini and Barba, 2006; Ilardi and Tavazza, 2015). Under experimental conditions, PPV can be transmitted mechanically to numerous <i>Prunus</i> spp. and several herbaceous plants e.g <i>Arabidopsis thaliana</i>, <i>Chenopodium foetidum</i>, <i>Nicotiana benthamiana</i>, <i>N. clevelandii</i>, <i>N. glutinosa</i> and <i>Pisum sativum</i>.</u>	Kenya Category : <i>TECHNICAL</i>	Modified (Changes made as recommended and general improvements made, such as an update of reference. Garcia and Cambra 2007 was changed to Garcia <i>et al.</i> , 2014.)

#	Para	Text	Comment	SC's response
28	54	<p><i>Plum pox virus</i> is a member of the genus <i>Potyvirus</i> in the family <i>Potyviridae</i>. The virus particles are flexuous rods of approximately 700 nm × 11 nm, and are composed of a single-stranded <u>positive sense</u> RNA molecule consisting of almost 10 000 nucleotides coated by up to 2 000 subunits of a single coat protein (García and Cambra, 2007). <u>PPV</u>; Ilardi and Tavazza, 2015). The genomic RNA carries a virus-encoded protein (VPg) at its 5' end and poly(A) tail at its 3' end. It possess a long open reading frame which is transmitted in the field by aphids in translated into a non-persistent manner large polyprotein precursor from the second AUG codon (García et al., but movement of infected propagative plant material 2014). The polyprotein is the main way in which later used to produce mature proteins. As an obligate intracellular pathogen with a limited genome capacity, PPV is spread over long distances unable to accomplish its infection cycle. Therefore it relies on the multifunction properties of its proteins and host factors for its infectivity process (García et al., 2014)</p> <p><u>Transmission:</u></p> <p><u>PPV is transmitted in the field by aphids in a non-persistent manner, but movement of infected propagative plant material is the main way in which PPV is spread over long distances.</u></p>	<p>Kenya</p> <p>Category : <i>SUBSTANTIVE</i></p>	<p>Modified</p> <p>(Section 1 was modified. Some changes were not made as recommended. Additional information on the genome organization and characteristics of the virus were not added as they were not seen as necessary or relevant for diagnostic purposes)</p>
29	55	<p><i>Plum pox virus</i> isolates can be classified currently into nine strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i>, 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i>, <i>P. domestica</i>, <i>P. persica</i> and <i>P. salicina</i>. EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European</p>	<p>Viet Nam</p> <p>para 63 move to before para 56</p> <p>Category : <i>EDITORIAL</i></p>	<p>Considered but not incorporated.</p> <p>(The existing format is consistent with other IPPC virus-related DPs such as that for Citrus tristeza virus. Also a section on host range has now been identified.)</p>

#	Para	Text	Comment	SC's response
		<p>countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i>, 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i>, 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i>, 2012). A novel sour cherry-adapted Tat strain, neither C nor CR, has also been proposed (Chirkov <i>et al.</i>, 2016).</p> <p><u>Under natural conditions, PPV readily infects fruit trees of the genus Prunus used as commercial varieties or rootstocks: <i>P. armeniaca</i>, <i>P. cerasifera</i>, <i>P. davidiana</i>, <i>P. domestica</i>, <i>P. mahaleb</i>, <i>P. marianna</i>, <i>P. mume</i>, <i>P. persica</i>, <i>P. salicina</i>, and interspecific hybrids between these species. <i>Prunus avium</i>, <i>P. cerasus</i> and <i>P. dulcis</i> may be infected occasionally. The virus also infects many wild and ornamental Prunus species such as <i>P. besseyi</i>, <i>P. cistena</i>, <i>P. glandulosa</i>, <i>P. insititia</i>, <i>P. laurocerasus</i>, <i>P. spinosa</i>, <i>P. tomentosa</i> and <i>P. triloba</i>. Under experimental conditions, PPV can be transmitted mechanically to numerous Prunus spp. and several herbaceous plants (<i>Arabidopsis thaliana</i>, <i>Chenopodium foetidum</i>, <i>Nicotiana benthamiana</i>, <i>N. clevelandii</i>, <i>N. glutinosa</i> and <i>Pisum sativum</i>).</u></p>		
30	55	<p><i>Plum pox virus</i> isolates can be classified currently into nine <u>monophyletic</u> strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i>, 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i>, <i>P. domestica</i>, <i>P. persica</i> and <i>P. salicina</i>. EA isolates are</p>	<p>European Union</p> <p>Category : TECHNICAL</p>	Incorporated.

#	Para	Text	Comment	SC's response
		geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain putative strain (Tat), neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).		
31	55	<i>Plum pox virus</i> isolates can be classified currently into nine strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i> , 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> . EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing <u>shows</u> an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a	Kenya <i>Category : EDITORIAL</i>	Incorporated (Except that 'show' is inserted instead of 'shows'.)

#	Para	Text	Comment	SC's response
		potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain, neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).		
32	55	<p><i>Plum pox virus</i><u>PPV Isolates/Strains</u></p> <p><i>Plum pox virus</i> isolates can be classified currently into nine strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i>, 2013;). <u>The strains have specific symptomatology, host range, epidemiology, pathogenicity, genome sequences and aphid transmissibility.</u> Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The <u>two</u> strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i>, <i>P. domestica</i>, <i>P. persica</i> and <i>P. salicina</i>. EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. <u>PPV-In European countries,PPV-C and PPV-CR isolates have been identified as the PPV strains infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified Whereas in several European countries. These isolates form two</u>Canada, a distinct strains that have been defined as PPV C and PPV CR. An atypical PPV strain PPV-A was detected in <i>P. domestica</i> in Canada (PPV-W) Thus representing a distinct PPV strain- PPV-W. This strain has since then been detected in several countries in Europe (James <i>et al.</i>, 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i>, 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i>, 2012). A novel sour cherry-adapted Tat strain, neither C nor CR, has also been proposed (Chirkov <i>et al.</i>, 2016).</p>	<p>Kenya</p> <p>Category : SUBSTANTIVE</p>	<p>Modified</p> <p>(Some changes made as recommended. Some changes recommended were not made as it would alter meaning e.g. "In European countries PPV-C and PPV CR ...etc..)</p>
33	55	<p><i>Plum pox virus</i> isolates can be classified currently into nine <u>monophyletic</u> strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An</p>	<p>EPPO</p> <p>Category : TECHNICAL</p>	<p>Incorporated</p>

#	Para	Text	Comment	SC's response
		(Ancestor Marcus) (James <i>et al.</i> , 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> . EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain putative strain (Tat), neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).		
34	55	<i>Plum pox virus</i> isolates can be classified currently into nine strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i> , 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> . EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since	Turkey <i>Category : TECHNICAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain putative strain (Tat), neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).		
35	55	<i>Plum pox virus</i> isolates can be classified currently into nine <u>monophyletic</u> strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor Marcus) (James <i>et al.</i> , 2013;). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> . EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain, neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).	Turkey <i>Category : TECHNICAL</i>	Incorporated
36	55	<i>Plum pox virus</i> isolates can be classified currently into nine strains: D (Dideron), M (Marcus), C (Cherry), EA (El Amar), W (Winona), Rec (Recombinant), T (Turkish), CR (Cherry Russian) and An (Ancestor	Philippines not appropriate <i>Category : EDITORIAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		Marcus) (James <i>et al.</i> , 2013;2013)). Most PPV isolates belong to the D and M strains. PPV D and M strains can easily infect <i>P. armeniaca</i> and <i>P. domestica</i> but differ in their ability to infect <i>P. persica</i> cultivars. The strains vary in their pathogenicity; for example, The M isolates generally cause faster epidemics and more severe symptoms than D isolates in <i>P. armeniaca</i> , <i>P. domestica</i> , <i>P. persica</i> and <i>P. salicina</i> . EA isolates are geographically restricted to Egypt and little information is available about their epidemiology and biological properties. PPV isolates infecting <i>P. avium</i> and <i>P. cerasus</i> have been identified in several European countries. These isolates form two distinct strains that have been defined as PPV-C and PPV-CR. An atypical PPV was detected in <i>P. domestica</i> in Canada (PPV-W) representing a distinct PPV strain. PPV W has since been detected in several countries in Europe (James <i>et al.</i> , 2013). In addition, natural recombinants between the D and M strains of PPV have been described as PPV-Rec, these showing an epidemiological behaviour similar to the D strain. A second type of recombinant strain has been reported in Turkey (T strain, Ulubaş as T strain (Ulubaş Sercçe <i>et al.</i> , 2009). A single isolate of PPV An has been described and it has been proposed as a potential ancestor of PPV M (Palmisano <i>et al.</i> , 2012). A novel sour cherry-adapted Tat strain, neither C nor CR, has also been proposed (Chirkov <i>et al.</i> , 2016).		
37	59	Synonym: Sharka virus	Viet Nam Which references to make this synonym? Category : EDITORIAL	Considered but not incorporated (There is no reference available, but Sharka virus is a recognized synonym by the scientific community)
38	62	3. Detection and Identification <u>Host range</u>	Kenya Category : TECHNICAL	Incorporated (A section on Host Range was created.)
39	63	Under natural conditions, PPV readily infects fruit trees of the genus <i>Prunus</i> used as commercial varieties or rootstocks: <i>P. armeniaca</i> , <i>P. cerasifera</i> , <i>P. davidiana</i> , <i>P. domestica</i> , <i>P. mahaleb</i> , <i>P. marianna</i> , <i>P. mume</i> , <i>P. persica</i> , <i>P. salicina</i> , and interspecific hybrids between these species. <i>Prunus avium</i> , <i>P. cerasus</i> and <i>P. dulcis</i> may be infected	Viet Nam para 63 move to before para 56 Category : EDITORIAL	Modified (It is included now in a section Host Range)

#	Para	Text	Comment	SC's response
		occasionally. The virus also infects many wild and ornamental <i>Prunus</i> species such as <i>P. besseyi</i> , <i>P. cistena</i> , <i>P. glandulosa</i> , <i>P. insititia</i> , <i>P. laurocerasus</i> , <i>P. spinosa</i> , <i>P. tomentosa</i> and <i>P. triloba</i> . Under experimental conditions, PPV can be transmitted mechanically to numerous <i>Prunus</i> spp. and several herbaceous plants (<i>Arabidopsis thaliana</i> , <i>Chenopodium foetidum</i> , <i>Nicotiana benthamiana</i> , <i>N. clevelandii</i> , <i>N. glutinosa</i> and <i>Pisum sativum</i>).		
40	63	Under natural conditions, PPV readily infects fruit trees of the genus <i>Prunus</i> used as commercial varieties or rootstocks: <i>P. armeniaca</i> , <i>P. cerasifera</i> , <i>P. davidiana</i> , <i>P. domestica</i> , <i>P. mahaleb</i> , <i>P. marianna</i> , <i>P. mume</i> , <i>P. persica</i> , <i>P. salicina</i> , and interspecific hybrids between these species. <i>Prunus avium</i> , <i>P. cerasus</i> and <i>P. dulcis</i> may be infected occasionally. The virus also infects many wild and ornamental <i>Prunus</i> species such as <i>P. besseyi</i> , <i>P. cistena</i> , <i>P. glandulosa</i> , <i>P. insititia</i> , <i>P. laurocerasus</i> , <i>P. spinosa</i> , <i>P. tomentosa</i> and <i>P. triloba</i> . Under experimental conditions, PPV can be transmitted mechanically to numerous <i>Prunus</i> spp. and several herbaceous plants (<i>Arabidopsis thaliana</i> , <i>Chenopodium foetidum</i> , <i>Nicotiana benthamiana</i> , <i>N. clevelandii</i> , <i>N. glutinosa</i> and <i>Pisum sativum</i>).	Kenya <i>Category : TECHNICAL</i>	Modified (A section 3.1 Host Range was created. Information on transmission has been moved to Section 1)
41	63	Under natural conditions, PPV readily infects fruit trees of the genus <i>Prunus</i> -(family Rosacea) used as commercial varieties or rootstocks: <u>Major hosts include</u> , <i>P. armeniaca</i> , <i>P. cerasifera</i> , <i>P. davidiana</i> , <i>P. domestica</i> , <i>P. mahaleb</i> , <i>P. marianna</i> , <i>P. mume</i> , <i>P. persica</i> , <i>P. salicina</i> , and interspecific hybrids between these species. <u>Occasionally</u> , <i>Prunus avium</i> , <i>P. cerasus</i> and <i>P. dulcis</i> may be infected occasionally. The virus also infects many wild and ornamental <i>Prunus</i> species such as <i>P. besseyi</i> , <i>P. cistena</i> , <i>P. glandulosa</i> , <i>P. insititia</i> , <i>P. laurocerasus</i> , <i>P. spinosa</i> , <i>P. tomentosa</i> and <i>P. triloba</i> . Under experimental conditions, PPV can be transmitted mechanically to numerous <i>Prunus</i> spp. and several herbaceous plants (<i>Arabidopsis thaliana</i> , <i>Chenopodium foetidum</i> , <i>Nicotiana benthamiana</i> , <i>N. clevelandii</i> , <i>N. glutinosa</i> and <i>Pisum sativum</i>).	Kenya <i>Category : TECHNICAL</i>	Modified (A section 3.1 Host Range was created. Information on transmission has been moved to section 1. Also improvements/updates made to information.)
42	64	Sharka <i>Plum pox virus</i> symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light-green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf symptoms are similar to those caused by	Viet Nam <i>Category : EDITORIAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		<p>other viruses, such as <i>American plum line pattern virus</i>. <i>Prunus cerasifera</i> cv. GF 31 shows rusty-brown corking and cracking of the bark. Flower symptoms can occur on petals (discoloration) of some <i>P. persica</i> cultivars when infected with PPV-M or in <i>P. glandulosa</i> infected with PPV-D. Infected fruits show chlorotic spots or lightly pigmented yellow rings or line patterns. Fruits may become deformed or irregular in shape and develop brown or necrotic areas under the discoloured rings. Some fruit deformations, especially in <i>P. armeniaca</i> and <i>P. domestica</i>, are similar to those caused by <i>Apple chlorotic leaf spot virus</i>. Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of <i>P. armeniaca</i> show typical pale rings or spots. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour. Symptom development and intensity depend strongly on the host plant and climatic conditions; for example the virus may be latent for several years in cold climates.</p>		
43	64	<p>Sharka symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light-green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf symptoms are similar to those caused by other viruses, such as <i>American plum line pattern virus</i>. <i>Prunus cerasifera</i> cv. GF 31 shows rusty-brown corking and cracking of the bark. Flower symptoms can occur on petals (discoloration) of some</p> <p><u>Flower symptoms can occur on petals (discoloration) of some <i>P. persica</i> cultivars when infected with PPV-M or in <i>P. glandulosa</i> infected with PPV-D. Infected fruits show chlorotic spots or lightly pigmented yellow rings or line patterns. Fruits may become deformed or irregular in shape and develop brown or necrotic areas under the discoloured rings. Some</u></p>	<p>Kenya</p> <p>Category : EDITORIAL</p>	<p>Modified (Symptoms consolidated under a new section 3.2 Symptoms)</p>

#	Para	Text	Comment	SC's response
		<p>fruit deformations, especially in <i>P. armeniaca</i> and <i>P. domestica</i>, are similar to those caused by <i>Apple chlorotic leaf spot virus</i>. Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of</p> <p><u>Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of <i>P. armeniaca</i> show typical pale rings or spots. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour. Symptom development and intensity depend strongly on the host plant and climatic conditions; for example the virus may be latent for several years in cold climates. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour.</u></p>		
44	64	<p>Sharka symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf symptoms are similar to those caused by other viruses, such as <u>disease symptoms</u></p>	<p>Kenya</p> <p>Category : EDITORIAL</p>	<p>Modified</p> <p>(Symptoms consolidated under a new section 3.2 Symptoms.)</p>

#	Para	Text	Comment	SC's response
		<p><u>Sharka symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light-green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf symptoms are similar to those caused by other viruses, such as American plum line pattern virus. <i>Prunus cerasifera</i> cv. GF 31 shows rusty-brown corking and cracking of the bark. Flower symptoms can occur on petals (discoloration) of some <i>P. persica</i> cultivars when infected with PPV-M or in <i>P. glandulosa</i> infected with PPV-D. Infected fruits show chlorotic spots or lightly pigmented yellow rings or line patterns. Fruits may become deformed or irregular in shape and develop brown or necrotic areas under the discoloured rings. Some fruit deformations, especially in <i>P. armeniaca</i> and <i>P. domestica</i>, are similar to those caused by Apple chlorotic leaf spot virus. Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of <i>P. armeniaca</i> show typical pale rings or spots. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour. Symptom development and intensity depend strongly on the host plant and climatic conditions; for example the virus may be latent for several years in cold climates.</u></p>		
45	64	<p>Sharka symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light-green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf symptoms are similar to those caused by other viruses, such as American plum line pattern virus. <i>Prunus cerasifera</i> cv. GF 31 shows rusty-brown corking and cracking of the bark. Flower symptoms can occur on petals (discoloration) of some <i>P. persica</i> cultivars when infected with PPV-M or in <i>P. glandulosa</i> infected with PPV-D. Infected fruits show chlorotic spots or lightly pigmented yellow rings or line patterns. Fruits may become deformed or irregular in shape and develop brown or necrotic areas under the discoloured rings. Some fruit deformations, especially in <i>P. armeniaca</i> and <i>P. domestica</i>, are similar to those caused by Apple</p>	<p>PPPO Need more explanation on flower symptoms occurring on petals. Not very clear. Category : EDITORIAL</p>	<p>Modified (More details of flower symptoms are provided. Also a reference is given. Symptoms consolidated under a new section 3.2 Symptoms)</p>

#	Para	Text	Comment	SC's response
		<i>chlorotic leaf spot virus</i> . Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of <i>P. armeniaca</i> show typical pale rings or spots. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour. Symptom development and intensity depend strongly on the host plant and climatic conditions; for example the virus may be latent for several years in cold climates.		
46	64	Sharka symptoms may appear on leaves, shoots, bark, petals, fruits and stones in the field. They are usually distinct on leaves early in the growing season and include mild light-green discoloration; chlorotic spots, bands or rings; vein clearing or yellowing; or leaf deformation. Some of these leaf-these symptoms are similar to those caused by other viruses, such as <i>American plum line pattern virus</i> -, <i>Prunus cerasifera</i> cv. GF 31 shows rusty-brown corking and cracking of the bark. Flower symptoms <u>discoloration</u> can occur on petals (discoloration) of some <i>P. persica</i> cultivars when infected with PPV-M or in <i>P. glandulosa</i> infected with PPV-D. Infected fruits show chlorotic spots or lightly pigmented yellow rings or line patterns. Fruits may become deformed or irregular in shape and develop brown or necrotic areas under the discoloured rings. Some fruit deformations, especially in <i>P. armeniaca</i> and <i>P. domestica</i> , are similar to those caused by <i>Apple chlorotic leaf spot virus</i> . Diseased fruits may show internal browning and gummosis of the flesh and reduced quality. In severe cases the diseased fruits drop prematurely from the tree. In general, the fruits of early maturing cultivars show more marked symptoms than those of late maturing cultivars. Stones from diseased fruits of <i>P. armeniaca</i> show typical pale rings or spots. The alcohol or spirits produced from diseased fruits are unmarketable owing to an undesirable flavour. Symptom development and intensity <u>severity</u> depend strongly on the host plant and climatic conditions; for example the . <u>The</u> virus may be latent for several years in cold climates.	Philippines <i>Category : SUBSTANTIVE</i>	Modified (More details of flower symptoms provided. Also a reference is given. Symptoms consolidated under a new section 3.2 Symptoms)
47	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account	European Union Remark: In summer shoots without leaves are shipped as grafting material, therefore only phloem tissue or buds can be used for testing.	Modified. A statement regarding dormant cuttings was added. This was included now in the Section 3.4

#	Para	Text	Comment	SC's response
		virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.	<i>Category : SUBSTANTIVE</i>	titled: Sampling for serological and molecular tests.
48	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples	European Union Reliability of the detection of PPV may depend on the method used. Molecular tools such as real-time RT-PCR should be able to detect low virus concentration even in young shoots. Suggestion is added in the paragraph. <i>Category : SUBSTANTIVE</i>	NOTED

#	Para	Text	Comment	SC's response
		can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		
49	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from <u>grafted</u> shoots that are at least one-year-old <u>one-year-old</u> and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. <u>Dormant buds can also be used in summer, especially for testing graftwood. RT-PCR or real time PCR should be used for detection of PPV in dormant buds and preferably also in mature leaves in summer, especially for detection of latent infections.</u> Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, shoots or branches, or complete spurs <u>spurs</u> , can be selected.	European Union SharCo : Containment of Sharka virus in view of EU-expansion, 2010. DE.3.2 : Procedures for sampling in nursery blocks and validated tools and protocols for reliable and accurate PPV detection. <i>Category : SUBSTANTIVE</i>	Modified (The paragraph was moved and modified. It is now in Section 3.4: Sampling for serological and molecular tests. Changes made as recommended, except where changes resulted in change of meaning.)
50	65	General guidance on Sampling for analysis	Kenya <i>Category : EDITORIAL</i>	Modified A Section 3.4: Sampling for serological and molecular tests has

#	Para	Text	Comment	SC's response
		<p><u>General guidance on sampling methodologies</u> is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.</p>		<p>been created that has all related sampling information.</p>
51	65	<p>General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves,</p>	<p>Ghana</p> <p>Category : <i>SUBSTANTIVE</i></p>	<p>Considered but not incorporated.</p> <p>(Since <i>Prunus</i> species, the host of PPV, are temperate crops, the concepts of dry season and rainy season are not as appropriate as spring and autumn.)</p>

#	Para	Text	Comment	SC's response
		collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime <u>springtime or during the rainy season</u> , samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn <u>autumn or dry or the hot climate</u> , mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter <u>winter or cold season</u> , dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		
52	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn <u>or dry season</u> are less reliable than tests done on samples collected earlier in the spring <u>spring or in the rainy season</u> . Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing	Ghana <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (Since Prunus species, the host of PPV, are temperate crops, the concepts of dry season and rainy season are not as appropriate as spring and autumn.)

#	Para	Text	Comment	SC's response
		houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		
53	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from <u>grafted</u> shoots that are at least one year old <u>one-year-old</u> and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. <u>Dormant buds can also be used in summer, especially for testing graftwood. RT-PCR or real time PCR should be used for detection of PPV in dormant buds and preferably also in mature leaves in summer, especially for detection of latent infections.</u> Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, shoots or branches, or complete spurs <u>spurs</u> , can be selected.	EPPO SharCo : Containment of Sharka virus in view of EU-expansion, 2010. DE.3.2 : Procedures for sampling in nursery blocks and validated tools and protocols for reliable and accurate PPV detection <i>Category : SUBSTANTIVE</i>	Modified (Sampling from dormant cuttings now described, and the recommendation for testing with RT-PCR or real-time PCR are indicated. Taking leaves from one year old shoots of a symptomless mature plant is different from taking leaves from one year old grafted shoots.)
54	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather	EPPO <i>Category : SUBSTANTIVE</i>	Modified (Sampling from dormant cuttings now described, and the recommendation for testing with RT-PCR or real-time PCR are

#	Para	Text	Comment	SC's response
		conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		indicated. Taking leaves from one year old shoots of a symptomless mature plant is different from taking leaves from one year old grafted shoots.)
55	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer	EPPO Remark: In summer shoots without leaves are shipped as grafting material, therefore only phloem tissue or buds can be used for testing. <i>Category : SUBSTANTIVE</i>	Modified (Sampling from dormant cuttings now described)

#	Para	Text	Comment	SC's response
		and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		
56	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old <u>one-year-old</u> and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.	Turkey <i>Category : EDITORIAL</i>	Considered but not incorporated ("taken from shoots that are at least one year old" is correct and more accurate.)
57	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). For field sampling, a appropriate <u>For field sampling, appropriate</u> sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the	Philippines <i>Category : EDITORIAL</i>	Modified A Section 3.4: Sampling for serological and molecular tests has been created that has all related sampling information. Also the wording has not been changed as sampling may occur in greenhouses or screen houses also)

#	Para	Text	Comment	SC's response
		growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis. Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.		
58	65	General guidance on sampling methodologies is provided in ISPM 31 (<i>Methodologies for sampling of consignments</i>). Appropriate sample selection is critical for PPV detection. Sampling should take into account virus biology and local climatic conditions, in particular the weather conditions during the growing season. If typical symptoms are present, samples should be collected of flowers, leaves or fruits showing symptoms. In symptomless plants, samples should be taken from shoots that are at least one year old and have mature or fully expanded leaves, collected from the middle of each of the main branches (detection is not reliable in shoots less than one year old). Samples should be collected from at least four different sites (e.g. four branches or four leaves) in each plant; this is critical because of the uneven distribution of PPV. Sampling should not be done during months with the highest temperatures. Tests on samples collected in the autumn are less reliable than tests done on samples collected earlier in the spring. Plant material should preferably be collected from the internal parts of the tree canopy. In springtime, samples can be flowers, shoots with fully expanded leaves, or fruits. In summer	Slovenia <i>Category : SUBSTANTIVE</i>	Modified (The paragraph was moved and modified. It is now in Section 3.4: Sampling for serological and molecular tests. Changes made as recommended, except where changes resulted in change of meaning.)

#	Para	Text	Comment	SC's response
		<p>and autumn, mature leaves and the skin of mature fruits collected from the field or packing houses can be used for analysis.</p> <p><u>Dormant buds can also be used in summer, especially for testing graftwood. RT-PCR or real time PCR must be used for detection of PPV in dormant buds and preferably also in mature leaves in summer, especially for detection of latent infections.</u></p> <p>Flowers, leaves, shoots and fruit skin can be stored at 4 °C for not more than 10 days before processing. Fruits can be stored for one month at 4 °C before processing. In winter, dormant buds or bark tissues from the basal part of twigs, shoots, or branches, or complete spurs can be selected.</p>		
59	66	<p>Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, <u>subsequent tests the confirmation should use serological methods and vice versa be performed preferably with a test with a higher analytical sensitivity that the one used for initial identification.. It is possible to confirm the results of a molecular test by another molecular test targeting a different genome region or sequence analysis.</u> Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.</p>	<p>European Union</p> <p><i>Category : SUBSTANTIVE</i></p>	Incorporated
60	66	<p>Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present</p>	<p>European Union</p> <p>SharCo : Containment of Sharka virus in view of EU-expansion, 2010. DE.3.2 : Procedures for sampling in nursery blocks and validated tools and protocols for reliable and accurate PPV detection.</p> <p><i>Category : SUBSTANTIVE</i></p>	Modified (It is now indicated that 'confirmation should be performed preferably with a test with a higher analytical sensitivity or if possible by a molecular test targeting a different genome region...')

#	Para	Text	Comment	SC's response
		or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, subsequent tests should use serological methods and vice versa. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.		
61	66	<p>Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection</p> <p><u>Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection</u></p> <p>organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, subsequent tests should use serological methods and vice versa. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.</p>	<p>Kenya</p> <p>Category : EDITORIAL</p>	<p>Incorporated</p> <p>(Reorganized as recommended.)</p>
62	66	<p>Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present</p>	<p>EPPO</p> <p>Category : SUBSTANTIVE</p>	<p>Incorporated</p> <p>(Reorganized as recommended. It is now indicated that 'confirmation should be performed preferably with a test with a higher analytical sensitivity or if possible by a molecular test targeting a different genome region...')</p>

#	Para	Text	Comment	SC's response
		or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, subsequent tests should use serological methods and vice versa. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.		
63	66	Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, <u>subsequent tests the confirmation should use serological methods and vice versa be performed preferably with a test with a higher analytical sensitivity than the one used for initial identification.</u> It is possible to confirm the results of a molecular test by another molecular test targeting a different genome region or sequence analysis. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.	EPPO The confirmation should preferably be done with a more sensitive test than the test used for the initial identification. <i>Category : SUBSTANTIVE</i>	Incorporated (Reorganized as recommended. It is now indicated that 'confirmation should be performed preferably with a test with a higher analytical sensitivity or if possible by a molecular test targeting a different genome region...
64	66	Detection of PPV can be achieved using a biological, serological or molecular test, test while identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. especially during routine diagnosis of a pest widely established in a country) country. In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method, subsequent tests should use serological methods and	Philippines <i>Category : EDITORIAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		vice versa. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.		
65	66	Detection of PPV can be achieved using a biological, serological or molecular test; identification requires either a serological or molecular test. A serological or molecular test is the minimum requirement to detect and identify PPV (e.g. during routine diagnosis of a pest widely established in a country). In instances where the national plant protection organization (NPPO) requires additional confidence in the identification of PPV (e.g. detection in an area where the virus is not known to be present or detection in a consignment originating in a country where the pest is declared to be absent), further tests may be done. Where the initial identification was done using a molecular method different, subsequent tests should use serological methods and vice versa preferably more sensitive method than the one used for initial identification. Further tests may also be done to identify the strain of PPV present. In all cases, positive and negative controls must be included in the tests. The recommended techniques are described in the following sections.	Slovenia Molecular methods for detection of PPV are more sensitive than serological ones therefore the result obtained by molecular method cannot always be confirmed by the use of serological analysis. <i>Category : TECHNICAL</i>	Incorporated (Reorganized as recommended. It is now indicated that 'confirmation should be performed preferably with a test with a higher analytical sensitivity or if possible by a molecular test targeting a different genome region...
66	68	In this diagnostic protocol, methods (including reference to brand names) are described as published, as these define the original level of sensitivity, specificity and reproducibility achieved. The use of names of reagents, chemicals or equipment in these diagnostic protocols implies no approval of them to the exclusion of others that may also be suitable. Laboratory procedures presented in the protocols may be adjusted to the standards of individual laboratories, provided that they are adequately validated.	European Union <i>Category : EDITORIAL</i>	NOTED The text in the main body of the document and the footnote has been adjusted to avoid repetition while still including all relevant information.
67	68	In this diagnostic protocol, methods (including reference to brand names) are described as published, as these define the original level of sensitivity, specificity and reproducibility achieved. The use of names of reagents, chemicals or equipment in these diagnostic protocols implies no approval of them to the exclusion of others that may also be suitable. Laboratory procedures presented in the protocols may be adjusted to the standards of individual laboratories, provided that they are adequately validated.	EPPO We note that this is an IPPC editorial policy <i>Category : EDITORIAL</i>	NOTED The text in the main body of the document and the footnote has been adjusted to avoid repetition while still including all relevant information.
68	68	In this diagnostic protocol, methods (including reference to brand names) are described as published, as these define the original level of sensitivity, specificity and reproducibility achieved. The use of names of reagents,	Uruguay Text deleted to avoid repetition with text in the footnote <i>Category : TECHNICAL</i>	Incorporated. The text in the main body of the document and the footnote has been adjusted to avoid repetition

#	Para	Text	Comment	SC's response
		chemicals or equipment in these diagnostic protocols implies no approval of them to the exclusion of others that may also be suitable. Laboratory procedures presented in the protocols may be adjusted to the standards of individual laboratories, provided that they are adequately validated.		while still including all relevant information.
69	74	For sample processing, approximately 0.2–0.5 g of fresh plant material is cut into small pieces and placed in a suitable tube or plastic bag. The sample is homogenized in approximately 4–10 ml (1:20 w/v) of extraction buffer using an electrical tissue homogenizer, or a manual roller, hammer or similar tool. The extraction buffer is phosphate-buffered saline (PBS) pH 7.2–7.4, containing 2% polyvinylpyrrolidone and 0.2% sodium diethyl dithiocarbamate (Cambra <i>et al.</i> , 1994), or an alternative suitably validated buffer. Plant material should be homogenized thoroughly and used fresh.	United States of America Second sentence: Or 1:10 as recommended by the ELISA kit manufacturer <i>Category : TECHNICAL</i>	Modified
70	74	For sample processing, approximately 0.2–0.5 g of fresh plant material is cut into small pieces and placed in a suitable tube or plastic bag. The sample is homogenized in approximately 4–10 ml (1:20 w/v) of extraction buffer using an electrical tissue homogenizer, or a manual roller, hammer or similar tool. The extraction buffer is phosphate-buffered saline (PBS) pH 7.2–7.4, containing 2% polyvinylpyrrolidone and 0.2% sodium diethyl dithiocarbamate (Cambra <i>et al.</i> , 1994), or an alternative suitably validated <u>suitable</u> buffer. Plant material should be homogenized thoroughly and used fresh.	European Union It is unclear what 'a validated buffer' means. <i>Category : TECHNICAL</i>	Incorporated
71	74	For sample processing, approximately 0.2–0.5 g of fresh plant material is cut into small pieces and placed in a suitable tube or plastic bag. The sample is homogenized in approximately 4–10 ml (1:20 w/v) of extraction buffer using an electrical tissue homogenizer, or a manual roller, hammer or similar tool. The extraction buffer is phosphate-buffered saline (PBS) pH 7.2–7.4, containing 2% polyvinylpyrrolidone and 0.2% sodium diethyl dithiocarbamate (Cambra <i>et al.</i> , 1994), or an alternative suitably validated buffer. Plant material should be homogenized thoroughly and used fresh.	EPPO It is unclear what 'a validated buffer' means <i>Category : TECHNICAL</i>	Incorporated
72	76	Double-antibody sandwich indirect enzyme-linked immunosorbent assay (DASI-ELISA), also called triple-antibody sandwich (TAS)-ELISA, should be performed according to Cambra <i>et al.</i> (1994) using a specific monoclonal antibody such as 5B-IVIA, following the manufacturer's instructions.	China DASI-ELISA and TAS-ELIS are different in their operating procedures. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (Double-antibody sandwich indirect enzyme-linked immunosorbent assay, DASI-ELISA, is similar to TAS-ELISA)
73	76	Double-antibody sandwich indirect enzyme-linked immunosorbent assay (DASI-ELISA), also called triple-antibody sandwich (TAS)-ELISA,	China The serological methods cannot be established based on monoclonal antibody 5B-IVIA. Most	Considered but not incorporated.

#	Para	Text	Comment	SC's response
		should be performed according to Cambra <i>et al.</i> (1994) using a specific monoclonal antibody such as 5B-IVIA, following the manufacturer's instructions antibody.	people do not have any clues how to get the Mabs. Category : <i>SUBSTANTIVE</i>	The test described was based on the use of the Mab 5B-IVIA. An NPPO may choose to use alternate tests and or reagents, as long as they are suitably validated.
74	77	The only monoclonal antibody currently demonstrated to detect all strains of PPV with high reliability, specificity and sensitivity is 5B-IVIA (Cambra <i>et al.</i> , 2006a). Optimal detection of isolates of strain CR requires adjustment of the extraction buffer to pH 6.0 (Chirkov <i>et al.</i> , 2013; Glasa <i>et al.</i> , 2013). In a DIAGPRO ¹ ring-test conducted by 17 laboratories using a panel of 10 samples, including both PPV-infected (PPV-D, PPV-M and PPV-D+M) and healthy samples from France and Spain, DASI-ELISA using the 5B-IVIA monoclonal antibody was 95% accurate (number of true negatives and true positives diagnosed by the technique, divided by the number of samples tested). This accuracy was greater than that achieved with either immunocapture reverse transcription-polymerase chain reaction (IC-RT-PCR) which was 82% accurate, or co-operational RT-PCR (Co-RT-PCR) which was 94% accurate (Olmos <i>et al.</i> , 2007; Cambra <i>et al.</i> , 2008). The proportion of true negatives (number of true negatives diagnosed by the technique, divided by the number of healthy plants) identified by DASI-ELISA using the 5B-IVIA monoclonal antibody was 99.0%, compared with real-time RT-PCR using purified nucleic acid (89.2%) or spotted samples (98.0%), or IC-RT-PCR (96.1%). Capote <i>et al.</i> (2009) also reported that there is a 98.8% probability that a positive result obtained in winter with DASI-ELISA using the 5B-IVIA monoclonal antibody was a true positive.	European Union Furthermore, it is desirable to include additional ELISA tests using other antisera. Many labs still prefer the use of polyclonal antisera, and it can be expected that these polyclonal antisera would be suitable for universal PPV detection as well. It is suggested to make an inventory of available antisera and their performance based on data provided by producers and/or available from tests performance studies and proficiency tests on representative isolates of all relevant strains. However, it should be noted that the quality of specific antisera might exhibit batch-to-batch variation and, therefore, needs verification of the performance. Category : <i>TECHNICAL</i>	Modified (Sentence added in Section 3.5.1 about batch to batch variation, the use of polyclonal antibodies and the need for proper validation of a test before use.)
75	77	The only This monoclonal antibody currently has demonstrated to detect all strains of PPV with high reliability, specificity and sensitivity is 5B-IVIA (Cambra <i>et al.</i> , 2006a). Optimal detection of isolates of strain CR requires adjustment of the extraction buffer to pH 6.0 (Chirkov <i>et al.</i> , 2013; Glasa <i>et al.</i> , 2013). In a DIAGPRO ¹ ring-test conducted by 17 laboratories using a panel of 10 samples, including both PPV-infected (PPV-D, PPV-M and PPV-D+M) and healthy samples from France and Spain, DASI-ELISA using the 5B-IVIA monoclonal antibody was 95% accurate (number of true negatives and true positives diagnosed by the technique, divided by the number of samples tested). This accuracy was greater than that achieved with either immunocapture reverse	Kenya Category : <i>EDITORIAL</i>	Considered but not incorporated. (The existing description is clearer and more accurate.)

#	Para	Text	Comment	SC's response
		transcription-polymerase chain reaction (IC-RT-PCR) which was 82% accurate, or co-operational RT-PCR (Co-RT-PCR) which was 94% accurate (Olmos <i>et al.</i> , 2007; Cambra <i>et al.</i> , 2008). The proportion of true negatives (number of true negatives diagnosed by the technique, divided by the number of healthy plants) identified by DASI-ELISA using the 5B-IVIA monoclonal antibody was 99.0%, compared with real-time RT-PCR using purified nucleic acid (89.2%) or spotted samples (98.0%), or IC-RT-PCR (96.1%). Capote <i>et al.</i> (2009) also reported that there is a 98.8% probability that a positive result obtained in winter with DASI-ELISA using the 5B-IVIA monoclonal antibody was a true positive.		
76	77	The only monoclonal antibody currently demonstrated to detect all strains of PPV with high reliability, specificity and sensitivity is 5B-IVIA (Cambra <i>et al.</i> , 2006a). Optimal detection of isolates of strain CR requires adjustment of the extraction buffer to pH 6.0 (Chirkov <i>et al.</i> , 2013; Glasa <i>et al.</i> , 2013). In a DIAGPRO ¹ ring-test conducted by 17 laboratories using a panel of 10 samples, including both PPV-infected (PPV-D, PPV-M and PPV-D+M) and healthy samples from France and Spain, DASI-ELISA using the 5B-IVIA monoclonal antibody was 95% accurate (number of true negatives and true positives diagnosed by the technique, divided by the number of samples tested). This accuracy was greater than that achieved with either immunocapture reverse transcription-polymerase chain reaction (IC-RT-PCR) which was 82% accurate, or co-operational RT-PCR (Co-RT-PCR) which was 94% accurate (Olmos <i>et al.</i> , 2007; Cambra <i>et al.</i> , 2008). The proportion of true negatives (number of true negatives diagnosed by the technique, divided by the number of healthy plants) identified by DASI-ELISA using the 5B-IVIA monoclonal antibody was 99.0%, compared with real-time RT-PCR using purified nucleic acid (89.2%) or spotted samples (98.0%), or IC-RT-PCR (96.1%). Capote <i>et al.</i> (2009) also reported that there is a 98.8% probability that a positive result obtained in winter with DASI-ELISA using the 5B-IVIA monoclonal antibody was a true positive.	EPPO <i>Category : TECHNICAL</i>	Modified (Sentence added in Section 3.5.1 about batch to batch variation, the use of polyclonal antibodies and the need for proper validation of a test before use.)
77	77	The only monoclonal antibody currently demonstrated to detect all strains of PPV with high reliability, specificity and sensitivity is 5B-IVIA (Cambra <i>et al.</i> , 2006a). Optimal detection of isolates of strain CR requires adjustment of the extraction buffer to pH 6.0 (Chirkov <i>et al.</i> , 2013; Glasa <i>et al.</i> , 2013). In a DIAGPRO ¹ ring-test conducted by 17 laboratories using	Philippines Would this protocol/procedure be available online for free once this Annex is approved <i>Category : SUBSTANTIVE</i>	Considered, but not incorporated (All adopted ISPMs and their annexes are publically available on the IPPC website. NPPOs and RPPOs need to be informed of availability)

#	Para	Text	Comment	SC's response
		a panel of 10 samples, including both PPV-infected (PPV-D, PPV-M and PPV-D+M) and healthy samples from France and Spain, DAS-ELISA using the 5B-IVIA monoclonal antibody was 95% accurate (number of true negatives and true positives diagnosed by the technique, divided by the number of samples tested). This accuracy was greater than that achieved with either immunocapture reverse transcription-polymerase chain reaction (IC-RT-PCR) which was 82% accurate, or co-operational RT-PCR (Co-RT-PCR) which was 94% accurate (Olmos <i>et al.</i> , 2007; Cambra <i>et al.</i> , 2008). The proportion of true negatives (number of true negatives diagnosed by the technique, divided by the number of healthy plants) identified by DAS-ELISA using the 5B-IVIA monoclonal antibody was 99.0%, compared with real-time RT-PCR using purified nucleic acid (89.2%) or spotted samples (98.0%), or IC-RT-PCR (96.1%). Capote <i>et al.</i> (2009) also reported that there is a 98.8% probability that a positive result obtained in winter with DAS-ELISA using the 5B-IVIA monoclonal antibody was a true positive.		online, and the existence of contact points to provide any assistance required. Procedures are provided with the commercial kits)
78	80	The conventional or biotin–streptavidin system of double-antibody sandwich (DAS)-ELISA should be performed using utilize kits based on the specific monoclonal antibody 5B-IVIA or on polyclonal antibodies that have been demonstrated to detect all strains of PPV without cross-reacting with other viruses or healthy plant material (Cambra <i>et al.</i> , 2006a; Capote <i>et al.</i> , 2009). The test should be done according to the manufacturer's instructions.	Philippines <i>Category : EDITORIAL</i>	Incorporated
79	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	United States of America There is a citation for two papers but these paper do not list Agdia's kit and just briefly mentioned the Bioreba kit/antibody for DAS ELISA. Both kits/antibodies have been successfully used for the PPV Eradication program in USA. The Agdia kit antibodies were developed by a Canadian scientist Dr. Ann Rochon. <i>Category : TECHNICAL</i>	Modified Kits and procedures other than those indicated in the DP may be used, as long as they are properly validated. It is difficult to cover all possible scenarios in this DP. The paragraph has been modified and appropriate references added to improve accuracy
80	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	European Union Regarding the second subsection, the referred papers do not provide the data substantiating the statements on the lack of sensitivity and specificity. <i>Category : TECHNICAL</i>	Modified The paragraph has been modified and appropriate references added to improve accuracy

#	Para	Text	Comment	SC's response
81	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	European Union http://www.bioreba.ch/popup.php?docFile=http://www.bioreba.ch/files/Product_Info/ELISA_Reagents/PPV_DAS_ELISA.pdf. <i>Category : SUBSTANTIVE</i>	Modified Kits and procedures other than those indicated in the DP may be used, as long as they are properly validated. It is difficult to cover all possible scenarios in this DP. Also the paragraph has been modified and appropriate references added to improve accuracy
82	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	EPPO <i>Category : SUBSTANTIVE</i>	Modified The paragraph has been modified and appropriate references added to improve accuracy
83	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	EPPO Regarding the second subsection, the referred papers do not provide the data substantiating the statements on the lack of sensitivity and specificity. <i>Category : TECHNICAL</i>	Modified The paragraph has been modified and appropriate references added to improve accuracy
84	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	Philippines If this will be adopted as a standard diagnostic protocol, recommended protocol should include only those with high sensitivity, reliability and accuracy to optimize resources available. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated Some basic tests required for a diagnostic result are indicated. Other tests may be used once properly validated
85	81	Whereas the 5B-IVIA monoclonal antibody detects all PPV strains specifically, sensitively and reliably, some polyclonal antibodies are not specific and have limited sensitivity (Cambra <i>et al.</i> , 1994; Cambra <i>et al.</i> , 2006a). The use of additional methods is therefore recommended in situations where polyclonal antibodies have been used in a test and the NPPO requires additional confidence in the identification of PPV.	Slovenia <i>Category : SUBSTANTIVE</i>	Modified The paragraph has been modified and appropriate references added to improve accuracy
86	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker	United States of America For clarity <i>Category : EDITORIAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		with-and less opportunity for prone to contamination (with the target DNA) than conventional PCR.		
87	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker with less opportunity for contamination (with the target DNA) than conventional PCR.	European Union The general introduction needs to be updated, since molecular tests are commonly used for routine and large-scale testing nowadays. Moreover, it is desirable to make an inventory of the currently used molecular tests, since the described tests have been designed long ago and new formats, such as LAMP, have been developed since then (e.g. Ion et al., 2016; https://ijair.org/administrator/components/com_jresearch/files/publications/36_IJAIR_1842_Final.pdf). It is desirable to provide data and/or references substantiating that all strains, including the recently described strains, will be detected by the respective tests. Furthermore, guidance on using RT-PCR and sequence analysis for detection and identification of PPV should be included. <i>Category : TECHNICAL</i>	Modified (The paragraph has been modified. Also a paragraph has been added to the introduction of Section 3: detection and identification, indicating the use of LAMP and NGS for PPV detection. These have not yet been fully validated. The DP describes basic tests for a diagnostic result. Other tests may be used once properly validated.)
88	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker with less opportunity for contamination (with the target DNA) than conventional PCR.	European Union Recently, researchers from Naktuinbouw (NL) designed a new real-time RT-PCR (Taqman) for universal PPV detection, based on currently available sequence data from GenBank. Preliminary results are promising and validation is in progress. If the performance characteristics fulfil the requirements, Naktuinbouw is willing to provide the details so that the test can be included in both the IPPC and EPPO protocols. <i>Category : TECHNICAL</i>	Considered but not incorporated. (The DP describes some basic tests for getting a diagnostic result. Other tests may be used once properly validated. This document is not meant to be a review article where all available tests are listed.)
89	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker with less opportunity for contamination (with the target DNA) than conventional PCR.	EPPO <i>Category : TECHNICAL</i>	Modified (The paragraph has been modified. Also a paragraph has been added to the introduction of the section detection and identification, indicating the use of LAMP and NGS for PPV detection. These have not yet been fully validated. The DP describes basic tests for a diagnostic result. Other tests may be used once properly validated.)
90	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological	EPPO The general introduction needs to be updated,	Modified

#	Para	Text	Comment	SC's response
		methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker with less opportunity for contamination (with the target DNA) than conventional PCR.	since molecular tests are commonly used for routine and large-scale testing nowadays. Moreover, it is desirable to make an inventory of the currently used molecular tests, since the described tests have been designed long ago and new formats, such as LAMP, have been developed since then (e.g. Ion et al., 2016; https://ijair.org/administrator/components/com_jresearch/files/publications/36_IJAIR_1842_Final.pdf). It is desirable to provide data and/or references substantiating that all strains, including the recently described strains, will be detected by the respective tests. Furthermore, guidance on using RT-PCR and sequence analysis for detection and identification of PPV should be included. <i>Category : TECHNICAL</i>	(The paragraph has been modified. Also a paragraph has been added to the introduction of the section detection and identification, indicating the use of LAMP and NGS for PPV detection. These have not yet been fully validated. The DP describes basic tests for a diagnostic result. Other tests may be used once properly validated.)
91	83	Molecular methods using reverse transcription-polymerase chain reaction (RT-PCR) may be more expensive or time consuming than serological methods, especially for large-scale testing. However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. The use of real-time RT-PCR also avoids the need for any post-amplification processing (e.g. gel electrophoresis) and is therefore quicker with less opportunity for contamination (with the target DNA) than conventional PCR.	Philippines However, molecular methods, especially real-time RT-PCR, are generally more sensitive than serological methods. <i>Category : SUBSTANTIVE</i>	Modified. To improve accuracy
92	84	With the exception of IC-RT-PCR (for which RNA isolation is not required), RNA extraction should be conducted using appropriately validated protocols. The samples should be placed in individual plastic bags to avoid cross-contamination during extraction. Alternatively, for real-time RT-PCR, spotted plant extracts, printed tissue sections or squashes of plant material can be immobilized on blotting paper or nylon membranes and analysed by real-time RT-PCR (Olmos <i>et al.</i> , 2005; Osman and Rowhani, 2006; Capote <i>et al.</i> , 2009). It is not recommended that spotted or tissue-printed samples be used in conventional PCR because of the lower sensitivity compared with real-time RT-PCR.	Philippines RNA extraction should be conducted using appropriately validated protocols. (Please include RNA Extraction Protocol as attachment of this Annex) <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. The paragraph was maintained as is for accuracy.
93	85	Each of the following methods describes the volume of extracted sample that should be used as a template. Depending on the sensitivity of the method, the minimum concentration of template required to detect PPV varies as follows: RT-PCR, 100 fg RNA template/ml; Co-RT-PCR, 1 fg RNA template/ml; and real-time RT-PCR, 2 fg RNA template/ml.	European Union The information does not seem very useful when testing field samples. It should be indicated, how much plant material is needed to obtain 100 fg of RNA. This will depend on the applied extraction protocols. However, if there are any data available to give an Information on the quantity of plant material it should be stated.	Considered but not incorporated. (Impossible to give accurate tissue volumes to obtain the RNA concentrations recommended as the concentration obtained after extraction will depend on method, tissue type, tissue quality, reagents, etc.)


#	Para	Text	Comment	SC's response
			<i>Category : TECHNICAL</i>	
94	85	Each of the following methods describes the volume of extracted sample that should be used as a template. Depending on the sensitivity of the method, the minimum concentration of template required to detect PPV varies as follows: RT-PCR, 100 fg RNA template/ml; Co-RT-PCR, 1 fg RNA template/ml; and real-time RT-PCR, 2 fg RNA template/ml.	EPPO <i>Category : TECHNICAL</i>	Considered but not incorporated. (Impossible to give accurate tissue volumes to obtain the RNA concentrations recommended as the concentration obtained after extraction will depend on method, tissue type, tissue quality, reagents, etc.)
95	85	Each of the following methods describes the volume of extracted sample that should be used as a template. Depending on the sensitivity of the method, the minimum concentration of template required to detect PPV varies as follows: RT-PCR, 100 fg RNA template/ml; Co-RT-PCR, 1 fg RNA template/ml; and real-time RT-PCR, 2 fg RNA template/ml.	Philippines We recommend deletion of this paragraph because we have to be certain on the sensitivity of the method if this will be adopted as standard diagnostic protocol. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (The paragraph provides useful and valuable information. Other tests and approaches may be used once properly validated.)
96	87	The RT-PCR primers used in this method are either the primers of Wetzel <i>et al.</i> (1991):	European Union Please provide more details on the performance of these tests, i.e. the sensitivity expressed in relative infection rate; specificity in terms of strains as provided in the paper of Wetzel <i>et al.</i> (1992) instead of isolates from different countries. Moreover, information on the sample preparation (RNA extraction) is not provided. <i>Category : TECHNICAL</i>	Modified. (A sentence was added and clarification provided. Sensitivity based on relative infection rates are not described in any available publication.)
97	87	The RT-PCR primers used in this method are either the primers of Wetzel <i>et al.</i> (1991):	EPPO Please provide more details on the performance of these tests, i.e. the sensitivity expressed in relative infection rate; specificity in terms of strains as provided in the paper of Wetzel <i>et al.</i> (1992) instead of isolates from different countries. Moreover, information on the sample preparation (RNA extraction) is not provided. <i>Category : TECHNICAL</i>	Modified. (A sentence was added and clarification provided. Sensitivity based on relative infection rates are not described in any available publication)
98	93	The 25 µl reaction mixture is composed as follows: 1 µM of each primer (P1 and P2, or the 3'NCR primer pair), 250 µM dNTPs, 1 unit <i>Avian myeloblastosis virus</i> (AMV) reverse transcriptase, 0.5 units Taq DNA polymerase, 2.5 µl 10× Taq polymerase buffer, 1.5 mM MgCl ₂ , 0.3% Triton X-100 and 5 µl RNA template. The reaction is performed under the following thermocycling conditions: 45 min at 42 °C, 2 min at 94 °C, 40 cycles of 30 s at 94 °C, 30 s at either 60 °C (P1 and P2 primers) or 62 °C (3'NCR primers), and 1 min at 72 °C, followed by a final extension for 10 min at 72 °C. The PCR products are analysed by gel	European Union In Austria 10 µl and a commercial one step kit is used. This may be considered in an additional PCR protocol. <i>Category : TECHNICAL</i>	Considered but not incorporated. The test is given as described in the publication. Alternative tests or modifications of existing tests may be used once properly validated.

#	Para	Text	Comment	SC's response
		electrophoresis. The P1/P2 pair of primers produces a 243 base pair (bp) amplicon and the 3'NCR primers produce a 220 bp amplicon.		
99	93	The 25 µl reaction mixture is composed as follows: 1 µM of each primer (P1 and P2, or the 3'NCR primer pair), 250 µM dNTPs, 1 unit <i>Avian myeloblastosis virus</i> (AMV) reverse transcriptase, 0.5 units Taq DNA polymerase, 2.5 µl 10× Taq polymerase buffer, 1.5 mM MgCl ₂ , 0.3% Triton X-100 and 5 µl RNA template. The reaction is performed under the following thermocycling conditions: 45 min at 42 °C, 2 min at 94 °C, 40 cycles of 30 s at 94 °C, 30 s at either 60 °C (P1 and P2 primers) or 62 °C (3'NCR primers), and 1 min at 72 °C, followed by a final extension for 10 min at 72 °C. The PCR products are analysed by gel electrophoresis. The P1/P2 pair of primers produces a 243 base pair (bp) amplicon and the 3'NCR primers produce a 220 bp amplicon.	EPPO In Austria 10 µl and a commercial one step kit is used. This may be considered in an additional PCR protocol <i>Category : TECHNICAL</i>	Considered but not incorporated. The test is given as described in the publication. Alternative tests or modifications of existing tests may be used once properly validated.
100	96	The immunocapture phase should be performed according to Wetzel <i>et al.</i> (1992), using plant sap extracted as in section 3.2 using individual tubes or plastic bags to avoid contamination.	European Union This test uses the same primers as the test described in 3.3.1. The difference concerns the sample preparation, i.e. immunocapture using not further specified polyclonal antibodies or 5B-IVIA. It should be noted that the antiserum of Sanofi, used for the original test described by Wetzel <i>et al.</i> (1992) is no longer available. Furthermore, immunocapture appears sensitive to cross contamination compared to direct RNA extraction (e.g.RNeasy), especially when positive samples are present. Finally, it is not clear if the pH of the extraction buffer should be adapted 6.0 (see 3.2.1). Therefore, we have doubts whether this test should be recommended. <i>Category : TECHNICAL</i>	Modified. In this DP it is indicated that other tests or reagents may be used, once properly validated. The comment in 3.2.1 (now 3.5.1) was in relation to the ELISA extraction buffer. The change of pH is needed for reliable detection of isolates of PPV Cherry Russian (CR) when using Mab 5B in ELISA. This has not been described as required for isolates of other strains using IC-RT PCR. IC-RT PCR is known to be more sensitive and it is normal laboratory practice to avoid cross contamination.
101	96	The immunocapture phase should be performed according to Wetzel <i>et al.</i> (1992), using plant sap extracted as in section 3.2 using individual tubes or plastic bags to avoid contamination.	EPPO This test uses the same primers as the test described in 3.3.1. The difference concerns the sample preparation, i.e. immunocapture using not further specified polyclonal antibodies or 5B-IVIA. It should be noted that the antiserum of Sanofi, used for the original test described by Wetzel <i>et al.</i> (1992) is no longer available. Furthermore, immunocapture appears sensitive to cross contamination compared to direct RNA extraction (e.g.RNeasy), especially when positive samples are present. Finally, it is not clear if the pH of the extraction buffer should be adapted 6.0 (see 3.2.1). Therefore, we have doubts whether this test should be recommended.	Modified. In this DP it is indicated that other tests or reagents may be used, once properly validated. The comment in 3.2.1 (3.5.1) was in relation to the ELISA extraction buffer. The change of pH is needed for reliable detection of isolates of PPV Cherry Russian (CR) when using Mab 5B-IVIA in ELISA. This has not been described as required for isolates of other strains using IC-RT PCR. IC-RT PCR is known to be more sensitive and it is normal laboratory practice to avoid cross contamination.

#	Para	Text	Comment	SC's response
			<i>Category : TECHNICAL</i>	
102	97	A dilution (1 µg/ml) is prepared of polyclonal antibodies or PPV-specific monoclonal antibody (5B-IVIA) in carbonate buffer pH 9.6. Aliquots of 100 µl diluted antibody are dispensed into PCR tubes and incubated at 37 °C for 3 h. The tubes are then washed twice with 150 µl sterile PBS-Tween (washing buffer), and rinsed twice with RNase-free water. Plant extract (100 µl; see section 3.2) is clarified by centrifugation (5 min at 15 500 g), and the supernatant added to the coated PCR tubes. The tubes are incubated for 2 h on ice or at 37 °C, and then washed three times with 150 µl sterile PBS-Tween. The RT-PCR reaction mixture is prepared as described in section 3.3.1 using the primers of Wetzel <i>et al.</i> (1992), and added directly to the coated PCR tubes. The amplification is performed as described in section 3.3.1.	Philippines It was mentioned that some Polyclonal Antibodies are not specific and have limited sensitivity, why recommend this? <i>Category : SUBSTANTIVE</i>	Modified. Optimization and validation is necessary before using any test. Some modification and clarification have been made to improve accuracy.
103	100	The RT-PCR primers used in this co-operational (Co)-RT-PCR are the primers of Olmos Wetzel <i>et al.</i> (2002)(1991):	European Union <i>Category : TECHNICAL</i>	Modified (Both Olmos et al., 2002 and Wetzel et al., 1992 primers are now indicated clearly.)
104	100	The RT-PCR primers used in this co-operational (Co)-RT-PCR are the primers of Olmos <i>et al.</i> (2002):	EPPO Is this still a commonly used method/test? No information is provided on the specificity. <i>Category : TECHNICAL</i>	Considered but not incorporated. The (Co)-RT-PCR is indicated as being 100 times more sensitive than RT-PCR so is a useful confirmational test, not necessarily a routine test.
105	109	Real-time RT-PCR can be performed using either TaqMan or SYBR® Green I. Two TaqMan methods have been described for universal detection of PPV (Schneider <i>et al.</i> , 2004; Olmos <i>et al.</i> , 2005). The primers and TaqMan probe used in the first method are those reported by Schneider <i>et al.</i> (2004):	European Union The test described by Varga and James (2005) presents another test for universal detection and identification of D and M strains. How does this test perform in comparison the previous one? The fact that it is a two-step protocol is a disadvantage because of the risk of cross contamination. Furthermore , the same remarks apply as for the previous tests. Since the nad5 primers have been designed by Menzel et al (2002), this reference should be included. <i>Category : TECHNICAL</i>	Modified. Descriptions of test comparisons have not yet been published. The Menzel et al. (2002) reference has been added.
106	109	Real-time RT-PCR can be performed using either TaqMan or SYBR® Green I. Two TaqMan methods have been described for universal detection of PPV (Schneider <i>et al.</i> , 2004; Olmos <i>et al.</i> , 2005). The primers and TaqMan probe used in the first method are those reported by Schneider <i>et al.</i> (2004):	EPPO <i>Category : TECHNICAL</i>	Modified. Descriptions of test comparisons have not yet been published. The Menzel et al. (2002) reference has been added.
107	109	<u>3.3.4.1</u> Real-time RT-PCR can be performed using either TaqMan or SYBR® Green I. Two TaqMan methods have been described for universal	Philippines	Considered but not incorporated.

#	Para	Text	Comment	SC's response
		detection of PPV (Schneider <i>et al.</i> , 2004; Olmos <i>et al.</i> , 2005). The primers and TaqMan probe used in the first method are those reported by Schneider <i>et al.</i> (2004):	<i>Category : EDITORIAL</i>	The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.
108	113	The 25 µl reaction mixture is composed as follows: 1× reaction mix (0.2 mM of each dNTP and 1.2 mM MgSO ₄), 200 nM each of forward and reverse primers, 100 nM TaqMan probe, 4.8 mM MgSO ₄ , 0.5 µl RT/Platinum® Taq mix (Superscript® One-Step RT-PCR with Platinum® Taq DNA polymerase; Invitrogen) ² and 5 µl RNA template. The RT-PCR is performed under the following thermocycling conditions: 15 min at 52 °C, 5 min at 95 °C, 60 cycles of 15 s at 95 °C, and 30 s at 60 °C. The PCR products are analysed in real-time according to the equipment manufacturer's instructions.	European Union In Austria 10 µl and a commercial one step kit is used. This may be considered in an additional PCR protocol. <i>Category : TECHNICAL</i>	Considered but not incorporated. The test is given as described in the publication. Alternative tests or modifications of existing tests may be used once properly validated.
109	113	The 25 µl reaction mixture is composed as follows: 1× reaction mix (0.2 mM of each dNTP and 1.2 mM MgSO ₄), 200 nM each of forward and reverse primers, 100 nM TaqMan probe, 4.8 mM MgSO ₄ , 0.5 µl RT/Platinum® Taq mix (Superscript® One-Step RT-PCR with Platinum® Taq DNA polymerase; Invitrogen) ² and 5 µl RNA template. The RT-PCR is performed under the following thermocycling conditions: 15 min at 52 °C, 5 min at 95 °C, 60 cycles of 15 s at 95 °C, and 30 s at 60 °C. The PCR products are analysed in real-time according to the equipment manufacturer's instructions.	EPPO In Austria 10 µl and a commercial one step kit is used. This may be considered in an additional PCR protocol <i>Category : TECHNICAL</i>	Considered but not incorporated. The test is given as described in the publication. Alternative tests or modifications of existing tests may be used once properly validated.
110	114	In this diagnostic protocol, methods (including reference to brand names) are described as published, as these define the original level of sensitivity, specificity and reproducibility achieved. The use of names of reagents, chemicals or equipment in these diagnostic protocols implies no approval of them to the exclusion of others that may also be suitable. Laboratory procedures presented in <u>This information is given for the protocols convenience of users of this protocol and does not constitute an endorsement by the CPM of the chemical, reagent and or equipment named. Equivalent products may be adjusted-used if they can be shown to lead to the standards of individual laboratories, provided that these are adequately validated same results.</u>	Uruguay Text deleted to avoid repetition with paragraph 68. Text added according to text agreed for footnotes <i>Category : TECHNICAL</i>	Incorporated. The text in the main body of the document and the footnote has been adjusted to avoid repetition while still including all relevant information.
111	116	<u>3.3.4.2</u> The primers and TaqMan probe used in the second method are those reported by Olmos <i>et al.</i> (2005):	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.

#	Para	Text	Comment	SC's response
112	123	3.3.4.3 Varga and James (2005) described a SYBR® Green I method for the simultaneous detection of PPV and identification of D and M strains:	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.
113	139	This section describes additional methods (using DASi-ELISA, RT-PCR, Co-RT-PCR and real-time RT-PCR) for identification of PPV strains (see Figure 1). Strain identification is not an essential component of PPV identification but an NPPO may wish to determine the identity of the strain to assist in predicting its epidemiological behaviour.	European Union It is confusing that this introduction refers to RT-PCR sequencing as the method for identification of strains, whereas only semi-specific molecular tests are described in this section. See also general comment. Furthermore, no information is included on the identification of the T, An and TAT strains mentioned in section 1. How should these strains be identified? Moreover, for sequence analysis, information should be provided on the regions to be used for identification as well as the database to be used for reference. <i>Category : TECHNICAL</i>	Modified. A sentence has been added to Paragraph 140 that indicates that no tests have been described for the identification of the strains An and T as as yet there is no publication describing such a test. Perhaps too few isolates of these strains have been described to allow validation.
114	139	This section describes additional methods (using DASi-ELISA, RT-PCR, Co-RT-PCR and real-time RT-PCR) for identification of PPV strains (see Figure 1). Strain identification is not an essential component of PPV identification but an NPPO may wish to determine the identity of the strain to assist in predicting its epidemiological behaviour.	EPPO It is confusing that this introduction refers to RT-PCR sequencing as the method for identification of strains, whereas only semi-specific molecular tests are described in this section. See also general comment. Furthermore, no information is included on the identification of the T, An and TAT strains mentioned in section 1. How should these strains be identified? Moreover, for sequence analysis, information should be provided on the regions to be used for identification as well as the database to be used for reference. <i>Category : TECHNICAL</i>	Modified. A sentence has been added to Paragraph 140 that indicates that no tests have been described for the identification of the strains An and T as as yet there is no publication describing such a test. Perhaps too few isolates of these strains have been described to allow validation.
115	139	This section describes additional methods <u>steps for identification of PPV Strains</u> (using DASi-ELISA, RT-PCR, Co-RT-PCR and real-time RT-PCR <u>RT-PCR</u>) for identification of PPV strains (see Figure 1). Strain identification is not an essential component of PPV identification but an NPPO may wish to determine the identity of the strain to assist in predicting its epidemiological behaviour.	Philippines <i>Category : SUBSTANTIVE</i>	Incorporated
116	141		United States of America Part of this table is confusing. For the box on the lower right, under "negative". Suggest it	Incorporated.

#	Para	Text	Comment	SC's response
			say instead "Plum pox virus present: atypical isolate of known strain (An, C, CR...) present, OR other undescribed strain" <i>Category : TECHNICAL</i>	
117	141	 Delete "atypical strain An, C, CR, D, EA, M, Rec, T or W present, or". <u>Suggest sequencing to confirm the first findings.</u>	China For PPV strains or other undescribed strains should be identified through sequencing and blasting to avoid false positive results. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated Guidance is provided in Section 4
118	141		Philippines Figure 1. Steps in the Identification of Strains of PPV <i>Category : SUBSTANTIVE</i>	Incorporated.
119	144	4.1 Serological identification of strains	China Firstly, it is very difficult for most people to get the Mabs. Secondly, sometimes, it is not very accurate when detecting with some antibodies, and maybe it confuses to determinate the results. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. Other tests and/or reagents may be used once they are properly validated.
120	145	DASI-ELISA for differentiation between the two main PPV strains (D and M) should be performed according to Cambra <i>et al.</i> (1994), using D- and M-specific monoclonal antibodies (Cambra <i>et al.</i>, 1994; Boscia <i>et al.</i>, 1997), according to the manufacturer's instructions.	European Union Consider removing this section, since the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
121	145	DASI-ELISA for differentiation between the two main PPV strains (D and M) should be performed according to Cambra <i>et al.</i> (1994), using D- and M-specific monoclonal antibodies (Cambra <i>et al.</i>, 1994; Boscia <i>et al.</i>, 1997), according to the manufacturer's instructions.	EPPO Consider removing this section, since it the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
122	145	DASI-ELISA for differentiation between the two main PPV strains (D and M) should be performed according to Cambra <i>et al.</i> (1994), using D- and M-specific monoclonal antibodies (Cambra <i>et al.</i>, 1994; Boscia <i>et al.</i>, 1997), according to the manufacturer's instructions.	China Firstly, it is very difficult for most people to get the Mabs. Secondly, sometimes, it is not very accurate when detecting with some antibodies, and maybe it confuses to determinate the results. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
123	145	DASI-ELISA for differentiation between the two main PPV strains (D and M) should be performed according to Cambra <i>et al.</i> (1994), using D- and	Philippines provide protocol as attachment to this Annex <i>Category : SUBSTANTIVE</i>	Considered but not incorporated All adopted ISPMs and their annexes are publically available on the IPPC website. NPPOs and RPPOs

#	Para	Text	Comment	SC's response
		M-specific monoclonal antibodies (Cambra <i>et al.</i> , 1994; Boscia <i>et al.</i> , 1997), according to the manufacturer's instructions.		need to be informed of availability online, and the existence of contact points to provide any assistance required. Procedures are provided with the commercial kits
124	146	This method has been validated in the DIAGPRO ring test, showing an accuracy of 84% for PPV-D detection and 89% for PPV-M detection (Olmos <i>et al.</i>, 2007; Cambra <i>et al.</i>, 2008). The 4D monoclonal antibody is PPV-D specific but does not react with all PPV-D isolates. Furthermore, the AL monoclonal antibody used for PPV-M detection reacts with isolates belonging to strains M, Rec and T because these groups share the same coat protein sequence. A molecular test is therefore required to differentiate between M, Rec and T strains detected using an M-specific monoclonal antibody.	European Union Consider removing this section, since the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
125	146	This method has been validated in the DIAGPRO ring test, showing an accuracy of 84% for PPV-D detection and 89% for PPV-M detection (Olmos <i>et al.</i>, 2007; Cambra <i>et al.</i>, 2008). The 4D monoclonal antibody is PPV-D specific but does not react with all PPV-D isolates. Furthermore, the AL monoclonal antibody used for PPV-M detection reacts with isolates belonging to strains M, Rec and T because these groups share the same coat protein sequence. A molecular test is therefore required to differentiate between M, Rec and T strains detected using an M-specific monoclonal antibody.	EPPO Consider removing this section, since it the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
126	146	This method has been validated in the DIAGPRO ring-test, showing an accuracy of 84% for PPV-D detection and 89% for PPV-M detection (Olmos <i>et al.</i> , 2007; Cambra <i>et al.</i> , 2008). The 4D monoclonal antibody is PPV-D specific but does not react with all PPV-D isolates. Furthermore, the AL monoclonal antibody used for PPV-M detection reacts with isolates belonging to strains M, Rec and T because these groups share the same coat protein sequence. A molecular test is therefore required to differentiate between M, Rec and T strains detected using an M-specific monoclonal antibody.	China Firstly, it is very difficult for most people to get the Mabs. Secondly, sometimes, it is not very accurate when detecting with some antibodies, and maybe it confuses to determinate the results. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
127	146	This method has been validated in the DIAGPRO ring test, showing an accuracy of 84% for PPV-D detection and 89% for PPV-M detection	China Firstly, it is very difficult for most people to get the Mabs. Secondly, sometimes, it is not very	Considered but not incorporated.

#	Para	Text	Comment	SC's response
		(Olmos <i>et al.</i>, 2007; Cambra <i>et al.</i>, 2008). The 4D monoclonal antibody is PPV-D specific but does not react with all PPV-D isolates. Furthermore, the AL monoclonal antibody used for PPV-M detection reacts with isolates belonging to strains M, Rec and T because these groups share the same coat protein sequence. A molecular test is therefore required to differentiate between M, Rec and T strains detected using an M-specific monoclonal antibody.	accurate when detecting with some antibodies, and maybe it confuses to determinate the results. <i>Category : SUBSTANTIVE</i>	(There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
128	147	Serological identification of PPV isolates from EA and C groups may be done by DAS-ELISA using the EA or the C specific monoclonal antibodies described by Myrta <i>et al.</i> (1998, 2000). However, these tests need to be validated.	European Union Consider removing this section, since the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
129	147	Serological identification of PPV isolates from EA and C groups may be done by DAS-ELISA using the EA or the C specific monoclonal antibodies described by Myrta <i>et al.</i> (1998, 2000). However, these tests need to be validated.	EPPO Consider removing this section, since it the described tests are already known not to detect certain isolates of target strains and are not validated. <i>Category : TECHNICAL</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
130	147	Serological identification of PPV isolates from EA and C groups may be done by DAS-ELISA using the EA or the C specific monoclonal antibodies described by Myrta <i>et al.</i> (1998, 2000). However, these tests need to be validated.	China Firstly, it is very difficult for most people to get the Mabs. Secondly, sometimes, it is not very accurate when detecting with some antibodies, and maybe it confuses to determinate the results. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (There is no published data that invalidates this test. Also, other tests and/or reagents may be used once they are properly validated.)
131	148	4.2 Molecular identification of strains	United States of America Listing all of these assays appears excessive. At the same time primers developed by Nemchinov et al for PPV-SoC detection are not listed. Nowadays people mostly use sequencing for this purposes, especially in cases of trade disputes. Even in developing countries conventional sequencing is not such a hurdle anymore. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. (Other tests and/or reagents may be used once they are properly validated.)
132	148	4.2 Molecular identification of strains	European Union Varga and James (2006): C, EA, W <i>Category : TECHNICAL</i>	Modified (Indicated in Section 4.2.4. Also the nad5 primers have the proper citation of Menzel et al. 2002)
133	148	4.2 Molecular identification of strains	EPPO Varga and James (2006): C, EA, W	Modified

#	Para	Text	Comment	SC's response
			<i>Category : TECHNICAL</i>	(Indicated in Section 4.2.4. Also the nad5 primers have the proper citation of Menzel et al. 2002)
134	149	4.2.1 Reverse transcription-polymerase chain reaction	European Union Add the sequence of the PM primer that is missing in section 4.2.1. Reverse transcription-polymerase chain reaction. The sequence of primer PM should be added just below the PD primer sequence. <i>Category : TECHNICAL</i>	Modified. (PM primer sequence is indicated in a separate line below the PD primer sequence.)
135	149	4.2.1 Reverse transcription-polymerase chain reaction	EPPO add the sequence of the PM primer that is missing in section 4.2.1. Reverse transcription-polymerase chain reaction. The sequence of primer PM should be added just below the PD primer sequence. <i>Category : TECHNICAL</i>	Modified. (PM primer sequence is indicated in a separate line below the PD primer sequence.)
136	150	4.2.1.1 PPV-D and PPV-M are identified using the primers described by Olmos <i>et al.</i> (1997):	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.
137	153	The 25 µl reaction mixture is composed as follows: 1 µM P1 primer, 1 µM of either PD or PM primer, 250 µM dNTPs, 1 unit AMV reverse transcriptase (10 units/µl), 0.5 units Taq DNA polymerase (5 units/µl), 2.5 µl 10× Taq polymerase buffer, 1.5 mM MgCl ₂ , 0.3% Triton X-100, 2% formamide and 5 µl RNA template. The RT-PCR is performed under the following thermocycling conditions: 45 min at 42 °C, 2 min at 94 °C, 40 <u>35</u> cycles of 30 s at 94 °C, 30 s at 60 °C, and 1 min <u>30 s</u> at 72 °C, followed by a final extension for 10 min at 72 °C. The PCR products are analysed by gel electrophoresis. The P1/PD pair of primers, and the P1/PM pair of primers, both produce a 198 bp amplicon. Olmos <i>et al.</i> (1997) evaluated their method using six isolates of PPV-D and four PPV-M isolates.	China 30 seconds is enough for extension when PCR is performed using the primer pairs (P1 and P2, or the 3'NCR primers). Also, we think 40 cycles are too many, 35 cycles are enough for PCR in general. <i>Category : TECHNICAL</i>	Considered but not incorporated. The test is described as published. Other tests and/or reagents may be used once they are properly validated.
138	154	4.2.1.2 The real-time reverse transcription-polymerase chain reaction with SYBR® Green I by Varga and James (2005) described in detail above in section 3.3.4 is also suitable for the identification of D and M strains of PPV.	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the

#	Para	Text	Comment	SC's response
				recommendation is to avoid a fourth level of section.
139	155	PPV-Rec is identified using the mD5 and mM3 Rec-specific primers described by Šubr <i>et al.</i> (2004):	European Union Missing PCR conditions. <i>Category : TECHNICAL</i>	Incorporated
140	155	PPV-Rec is identified using the mD5 and mM3 Rec-specific primers described by Šubr <i>et al.</i> (2004):	EPPO Missing PCR conditions <i>Category : TECHNICAL</i>	Incorporated
141	155	<u>4.2.1.3</u> PPV-Rec is identified using the mD5 and mM3 Rec-specific primers described by Šubr <i>et al.</i> (2004):	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. (see response to comment 107)
142	159	<u>4.2.1.4</u> PPV-CR is identified using the CR8597F and CR9023R primers described by Glasa <i>et al.</i> (2013):	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. (see response to comment 107)
143	162	A two-step RT-PCR protocol is used for specific detection of PPV-CR isolates (Glasa <i>et al.</i> , 2013). Complementary (c)DNA is synthesized from total RNA extracts (NucleoSpin® RNA Plant Kit, Macherey-Nagel ¹²) using random hexamer primers and AMV reverse transcriptase. An aliquot of cDNA is then added to the PCR reaction mix containing EmeraldAmp GT PCR Master Mix (TaKaRa Bio Inc. ¹²). The PCR is performed under the following thermocycling conditions: 1 min at 98 °C, 35 cycles of 98 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s, followed by a final extension at 72 °C for 5 min. The PCR products are analysed by gel electrophoresis. The CR-specific primers amplify a product 427 bp in size, targeting the 5' terminal CP coding region. The specificity of the CR primers was validated using isolates of PPV strains D, M, Rec, T, W, EA and C (Glasa <i>et al.</i> , 2013).	European Union <i>Category : EDITORIAL</i>	Incorporated
144	162	A two-step RT-PCR protocol is used for specific detection of PPV-CR isolates (Glasa <i>et al.</i> , 2013). Complementary (c)DNA is synthesized from total RNA extracts (NucleoSpin® RNA Plant Kit, Macherey-Nagel ¹) using random hexamer primers and AMV reverse transcriptase. An aliquot of cDNA is then added to the PCR reaction mix containing EmeraldAmp GT PCR Master Mix (TaKaRa Bio Inc. ¹). The PCR is performed under the following thermocycling conditions: 1 min at 98 °C, 35 cycles of 98 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s, followed by a final extension at 72 °C for 5 min. The PCR products are analysed by gel electrophoresis. The CR-specific primers amplify a product 427 bp in size, targeting the 5' terminal CP coding region. The specificity of the CR primers was	Japan Add information on master mix composition for two-step RT-PCR. It is necessary information in molecular methods. <i>Category : TECHNICAL</i>	Considered but not incorporated. The EmeraldAmp GT PCR Master Mix (TaKaRa Bio) is proprietary information. Other tests and/or reagents may be used once they are properly validated

#	Para	Text	Comment	SC's response
		validated using isolates of PPV strains D, M, Rec, T, W, EA and C (Glasa <i>et al.</i> , 2013).		
145	162	A two-step RT-PCR protocol is used for specific detection of PPV-CR isolates (Glasa <i>et al.</i> , 2013). Complementary (c)DNA is synthesized from total RNA extracts (NucleoSpin® RNA Plant Kit, Macherey-Nagel ⁴²) using random hexamer primers and AMV reverse transcriptase. An aliquot of cDNA is then added to the PCR reaction mix containing EmeraldAmp GT PCR Master Mix (TaKaRa Bio Inc. ⁴³). The PCR is performed under the following thermocycling conditions: 1 min at 98 °C, 35 cycles of 98 °C for 30 s, 55 °C for 30 s, 72 °C for 30 s, followed by a final extension at 72 °C for 5 min. The PCR products are analysed by gel electrophoresis. The CR-specific primers amplify a product 427 bp in size, targeting the 5' terminal CP coding region. The specificity of the CR primers was validated using isolates of PPV strains D, M, Rec, T, W, EA and C (Glasa <i>et al.</i> , 2013).	EPPO We note that this is IPPC editorial policy. <i>Category : EDITORIAL</i>	NOTED
146	171	<u>4.2.4.1</u> PPV-D and PPV-M are specifically identified using either SYBR® Green I chemistry according to the method of Varga and James (2005) (see section 3.3.4) or the TaqMan method described by Capote <i>et al.</i> (2006).	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.
147	178	<u>4.2.4.2</u> PPV-C, PPV-EA and PPV-W are specifically identified using SYBR® Green I chemistry according to the method of Varga and James (2006). The primers used in this method are:	Philippines <i>Category : EDITORIAL</i>	Considered but not incorporated. The draft DP has been edited and modified and the section numbers changed. Even though it is possible to have a four level section, the recommendation is to avoid a fourth level of section.
148	179	P1 (5' ACC GAG ACC ACT ACA CTC CC 3')	China One-step RT-SYBR Green was used for the detection of PPV-C, -EA, -W but P1 primer was not used here, and I suggest deleting the P1 primer. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. The P1 primer is an essential part of the one-tube real-time RT-PCR described by Varga and James (2006a).
149	192	Positive nucleic acid control. This control is used to monitor the efficiency of the test method (apart from the extraction) and, in RT-PCR, the amplification. Pre-prepared (stored) Total plant or viral RNA or PPV-infected plant material material, including printed on a membrane may be	United States of America Plasmid DNA controls also could be used if PPV infected tissue or total plant/virus RNA is not available. <i>Category : TECHNICAL</i>	Modified. PPV-infected plant material printed on a membrane is more accurate in this context so kept.

#	Para	Text	Comment	SC's response
		used. The stored RNA or PPV preparations should be verified periodically to determine the quality of the control with increased storage time.		
150	192	Positive nucleic acid control. This control is used to monitor the efficiency of the test method (apart from the extraction) and, in RT-PCR, the amplification. Pre-prepared (stored) RNA or PPV-infected plant material printed on a membrane may be used. The stored RNA or PPV preparations should be verified periodically to determine the quality of the control with increased storage time-time <u>plus d'éclaircissement sur l'utilité de l'emploi du terme healthy plant (plant sain) dans le cas du positive extraction control sachant que ce dernier ne peut pas être extrait d'un healthy plant ;</u>	Algeria <i>Category : TECHNICAL</i>	Considered but not incorporated. Negative controls are described separately.
151	192	Positive nucleic acid control. This control is used to monitor the efficiency of the test method (apart from the extraction) and, in RT-PCR, the amplification. Pre-prepared (stored) RNA or PPV-infected plant material printed on a membrane may be used. The stored RNA or PPV preparations should be verified periodically to determine the quality of the control with increased storage time.	Philippines we suggest that this para will recommend the use of Certified Reference Material, which is important in case of dispute in diagnosis arise bet NPPO. <i>Category : SUBSTANTIVE</i>	Considered but not incorporated. If available a NPPO may certainly decide to use certified reference material.
152	193	Internal control. For the real-time RT-PCR, mRNA of the mitochondrial gene <i>NADH dehydrogenase 5 (nad5)</i> could be incorporated into the RT-PCR protocol as an internal control to eliminate the possibility of RT-PCR false negatives due to nucleic acid extraction failure or degradation or the presence of RT-PCR inhibitors.	United States of America Menzel, W., Jelkmann, W., Maiss, E., 2002. Detection of four apple viruses by multiplex RT-PCR assays with coamplification of plant mRNA as internal control. J. Virol. Methods 99, 81–92. <i>Category : TECHNICAL</i>	Incorporated.
153	195	Positive extraction control. This control is used to ensure that the target nucleic acid extracted is of sufficient quantity and quality for RT-PCR and that the target virus is detectable. Nucleic acid is extracted from <u>PPV</u> infected host tissue, or healthy plant or insect tissues that have been spiked with PPV.	United States of America For clarity <i>Category : EDITORIAL</i>	Incorporated
154	197	Negative extraction control. This control is used to monitor contamination during nucleic acid extraction and cross-reaction with the host tissue. The control comprises nucleic acid that is extracted from uninfected host tissue and subsequently amplified. It is recommended that multiple controls be included <u>in random order</u> when large numbers of positive samples are expected.	United States of America For accuracy <i>Category : TECHNICAL</i>	Incorporated
155	198	In the case of immunocapture RT-PCR where no nucleic extraction is performed, plant sap from a known PPV positive should be used as a positive control, and plant sap from a healthy plant should be used as a negative control. A negative amplification control may also be included. The latter control is used to rule out	United States of America Negative amplification control described in 194 still may be needed. <i>Category : TECHNICAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		false positives due to contamination during the preparation of the reaction mixture. RNase-free PCR-grade water that was used to prepare the reaction mixture is added at the amplification stage for use as a negative amplification control.		
156	201	In instances where other contracting parties may be affected by the results of the diagnosis, in particular in cases of non-compliance and where the virus is found in an area for the first time, the following additional material should be kept:	United States of America This section should give some guidance on duration of time to keep/preserve the samples. As mentioned, different labs follow different protocols. <i>Category : TECHNICAL</i>	Incorporated
157	201	In instances where other contracting parties may be affected by the results of the diagnosis, in particular in cases of non-compliance and where the virus is found in an area for the first time, the following additional material should be kept kept for at least one year:	European Union Important precision to be given (duration consistent with other DPs). <i>Category : SUBSTANTIVE</i>	Incorporated
158	201	In instances where other contracting parties may be affected by the results of the diagnosis, in particular in cases of non-compliance and where the virus is found in an area for the first time, the following additional material should be kept kept for at least one year:	EPPO Important precision to be given (duration consistent with other DPs). <i>Category : SUBSTANTIVE</i>	Incorporated
159	202	The original sample (labelled appropriately for traceability) should be kept frozen-frozen, if possible, at – 80 °C or freeze-dried and kept at room temperature.	United States of America See United States comment in paragraph 201 <i>Category : TECHNICAL</i>	Incorporated
160	203	If relevant, RNA extractions-extracts should be kept at – 80 °C and spotted plant extracts or printed tissue sections (paper on paper or nylon membranes) should be kept at room temperature.	European Union <i>Category : EDITORIAL</i>	Incorporated
161	203	If relevant, RNA extractions-extracts should be kept at – 80 °C and spotted plant extracts or printed tissue sections (paper on paper or nylon membranes) should be kept at room temperature.	EPPO <i>Category : EDITORIAL</i>	Incorporated
162	205	6. Contact Points for Further Information	Viet Nam This section move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
163	206	United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), Plant Protection and Quarantine (PPQ), Registrations Identifications Permits and Plant Safeguarding	Viet Nam para 206 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.

#	Para	Text	Comment	SC's response
		(RIPPS), Molecular Diagnostic Laboratory, BARC Building 580, Powder Mill Road, Beltsville, Maryland 20705, United States of America (Ms Laurene Levy, e-mail: Laurene.Levy@aphis.usda.gov ; tel.: +1 3015045700; fax: +1 3015046124).		
164	206	United States Department of Agriculture (USDA), Animal and Plant Health Inspection Service (APHIS), Plant Protection and Quarantine (PPQ), Registrations Identifications Permits Science and Plant Safeguarding (RIPPS), Molecular Diagnostic Technology Beltsville Laboratory, BARC Building Bldg. 580, BARC-East, Powder Mill Road Rd., Beltsville, Maryland MD 20705, United States of America-USA (Ms Laurene Levy Vessela Mavrodieva, e-mail: vessela.a.mavrodieva@aphis.usda.gov , tel: +1 3013139208; fax: +1 3023139232). Laurene.Levy@aphis.usda.gov ; tel.: +1 3015045700; fax: +1 3015046124).	United States of America Updated the USDA laboratory contact information. <i>Category : EDITORIAL</i>	Incorporated.
165	207	Equipe de Virologie Institut National de la Recherche Agronomique (INRA), Centre de Bordeaux, UMR GD2P, IBVM, BP 81, F 33883 Villenave d'Ornon Cedex, France (Mr Thierry Candresse, e-mail: tc@bordeaux.inra.fr ; tel.: +33 557122389; fax: +33 557122384).	Viet Nam para 207 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
166	208	Faculty of Horticultural Science, Department of Plant Pathology, Corvinus University, Villányi út 29 43, H 1118 Budapest, Hungary (Mr Laszlo Palkovics, e-mail: laszlo.palkovics@uni-corvinus.hu ; tel.: +36 14825438; fax: +36 14825023).	Viet Nam para 208 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
167	209	Institute of Virology, Slovak Academy of Sciences, Dúbravská, 84505 Bratislava, Slovakia (Mr Miroslav Glasa, e-mail: virumig@savba.sk ; tel.: +421 259302447; fax: +421 254774284).	Viet Nam para 2096 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
168	210	Instituto Valenciano de Investigaciones Agrarias (IVIA), Plant Protection and Biotechnology Centre, Carretera Moncada Náquera km 5, 46113 Moncada (Valencia), Spain (Mr Antonio Olmos, e-mail: aolmos@ivia.es ; tel.: +34 963424000; fax: +34 963424001).	Viet Nam para 210 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
169	211	Istituto di Virologia Vegetale del CNR, sezione di Bari, via Amendola 165/A, I 70126 Bari, Italy (Mr Donato Boscia, e-mail: d.boscia@ba.ivv.cnr.it ; tel.: +39 0805443067; fax: +39 0805442911).	Viet Nam para 211 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
170	212	Sidney Laboratory, Canadian Food Inspection Agency (CFIA), British Columbia, V8L 1H3 Sidney, Canada (Mr Delano James, e-mail: Delano.James@inspection.gc.ca ; tel.: +1 250 3636650; fax: +1 250 3636661).	Viet Nam para 212 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.

#	Para	Text	Comment	SC's response
171	213	A request for a revision to a diagnostic protocol may be submitted by national plant protection organizations (NPPOs), regional plant protection organizations (RPPOs) or Commission on Phytosanitary Measures (CPM) subsidiary bodies through the IPPC Secretariat (ippc@fao.org), who will in turn forward it to the Technical Panel on Diagnostic Protocols (TPDP).	Viet Nam para 213 move to Appendix 1 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
172	214	7. Acknowledgements	Viet Nam This section move to Appendix 2 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
173	215	This diagnostic protocol was drafted by Mr M. Cambra, Mr A. Olmos and N. Capote, IVIA (see preceding section); Mr N.L. Africander, Department of Agriculture, Forestry and Fisheries, Private Bag X 5015, Stellenbosch, 75999, South Africa; Ms L. Levy, USDA, United States of America (see preceding section); Mr S.L. Lenardon, Instituto de Fitopatología y Fisiología Vegetal Instituto Nacional de Tecnología Agropecuaria (IFFIVE INTA), Cno. 60 Cuadras Km 51/2, Córdoba X5020ICA, Argentina; Mr G. Clover, Plant Health & Environment Laboratory, Ministry of Agriculture and Forestry, PO Box 2095, Auckland 1140, New Zealand; and Ms D. Wright, Plant Health Group, Central Science Laboratory, Sand Hutton, York YO41 1LZ, United Kingdom.	Viet Nam para 215 move to Appendix 2 <i>Category : EDITORIAL</i>	Considered but not incorporated. The current format is in line with the IPPC protocol's format.
174	235	García, J.A. & Cambra, M. 2007. Plum pox virus and sharka disease. <i>Plant Viruses</i>, 1: 69–79. <u>García, J.A., Glasa, M., Cambra, M., and Candresse, T. (2014). Plum pox virus and sharka: a model potyvirus and a major disease. <i>Mol. Plant Pathol.</i> 15, 226–241. doi:10.1111/mpp.12083</u>	Kenya <i>Category : TECHNICAL</i>	Incorporated.
175	238	<u>James Ilardi, V & Tarazza, D., Varga, A.M. & Sanderson 2015. Biotechnological strategies and tools for plum pox resistance; trans-, intra-, & cis-genesis and beyond. <i>Frontiers in plant science</i>, 6:1-16</u>	Kenya <i>Category : TECHNICAL</i>	Incorporated

#	Para	Text	Comment	SC's response
		James, D., Varga, A. & Sanderson, D. 2013. Genetic diversity of <i>Plum pox virus</i> : Strains, disease and related challenges for control. <i>Canadian Journal of Plant Pathology</i> , 35: 431–441.		
176	240	Levy, L. & Hadidi, A. 1994. A simple and rapid method for processing tissue infected with <i>Plum pox potyvirus</i> for use with specific 3' non-coding region RT-PCR assays. <i>EPPO Bulletin</i> , 24: 595–604. <u>Menzel W, Jelkmann W & Maiss E. 2002. Detection of four apple viruses by multiplex RT-PCR tests with co-amplification of plant mRNA as internal control. <i>Journal of Virological Methods</i> 99, 81–92.</u>	European Union Add the following reference (see comment on section 3.3.4). <i>Category : TECHNICAL</i>	Incorporated
177	240	Levy, L. & Hadidi, A. 1994. A simple and rapid method for processing tissue infected with <i>Plum pox potyvirus</i> for use with specific 3' non-coding region RT-PCR assays. <i>EPPO Bulletin</i> , 24: 595–604. <u>Menzel W, Jelkmann W & Maiss E (2002) Detection of four apple viruses by multiplex RT-PCR tests with co-amplification of plant mRNA as internal control. <i>Journal of Virological Methods</i> 99, 81–92.</u>	EPPO add the following reference (see comment on section 3.3.4) <i>Category : TECHNICAL</i>	Incorporated
178	251	Šubr, Z., Pittnerova, S. & Glasa, M. 2004. A simplified RT-PCR-based detection of recombinant <i>Plum pox virus</i> isolates. <i>Acta Virologica</i> , 48: 173–176. <u>Teshale, 2014. Evaluation of Molecular and Serological diagnostic techniques for a large scale detection of plum pox virus. <i>Research in Plant sciences</i>, 2:33-41</u>	Kenya <i>Category : TECHNICAL</i>	Considered but not incorporated. Not clear where this reference should be cited in the text.

