The European and Mediterranean Plant Protection Organization

EPPO activities

Nico Horn
Director-General



20 members from 17 countries plus European Commission



Scope

The Panel circulates information and develops harmonised views and positions for NPPOs on all matters relating to:

- the IPPC,
- the Commission on Phytosanitary Measures,
- relations with other RPPOs
- and other relevant international agreements (e.g. the Sanitary and Phytosanitary Agreement).

Tasks

- Develops a common EPPO view on strategic CPM issues
- Agreeing EPPO comments on draft Standards (ISPMs)
- Loading comments on OCS for EPPO member countries to endorse
- Reviewing documents in preparation for CPM meetings
- Coordinate Call for Topics for new Standards and Implementation material

Tasks

- Propose nominations for Europe region representatives on CPM bodies (for endorsement by EPPO Council)
- Capacity building for input into draft ISPMs
- Capacity building for implementation of ISPMs
- Collaboration with RPPOs (e.g. TC, coordination meetings)
- Finding European stewards for draft ISPMs

Way of working

- It is expected that experts from the EPPO region serving on CPM Bodies will also be members of the Panel
- The Panel normally meets three times per year (use of teleconference)
- In between regular meetings teleconferences or e-mail consultations when needed.

IPPC Regional Workshops

The objectives of IPPC Regional Workshops are:

- analyze and prepare comments on draft ISPMs;
- build phytosanitary capacity and raise awareness on various activities of the IPPC Community; and
- provide a forum to exchanging experiences and ideas at the regional level.
- Input for the Panel discussion to formulate EPPO comments and position


IPPC RW Europe and Central Asia (ECA)







2. EPPO Expert Working Groups for PRA




Communication on PRA

PRA documents available in EPPO Global Database and

Platform on PRAs:


Datasheets Reports of PRA



These working-procedures provide to EPPO member countries appropriate information for the technical justification of phytosanitary measures established for certain pests


3. Guidelines on the design and implementation of a buffer zone

- Scope: provides general guidance on how to design buffer zones to minimize the probability of spread of a pest into or out of delimited areas
- Public: risk assessors and risk managers, to help traceability
 of recommendations in Pest Risk Analysis or contingency
 plans, or when deciding on measures for an outbreak



Key element: Estimation of dispersal behaviour

- Based on Literature review, data retrieval & expert judgement
- Dispersal behaviour usually described by a dispersal kernel



- When possible, models analyzing suitable data should be used
- Estimation of a risk parameter can also be done using expert knowledge elicitation (EKE): estimation of the range, median, lower and upper quartile of a parameter
- -> EWG with experts on the biology, risk manager & a facilitator

Trirachys sartus (AELSSA)

4. Revision of EPPO Datasheets (±400)

MENU

Overview
Distribution
Host plants
Host commodities
Categorization
Reporting
Documents
Datasheet →

TOOLS

Save as word file
Save as PDF file

EPPO Datasheet: Trirachys sartus

Last updated: 2020-06-10

IDENTITY

Preferred name: Trirachys sartus

Authority: (Solsky)

Taxonomic position: Animalia: Arthropoda: Hexapoda: Insecta: Coleoptera: Cerambycidae

Other scientific names: Aeolesthes sarta (Solsky), Pachydissus sartus Solsky

Common names in English: Uzbek longhorn beetle, city longhorn beetle, sart longhorn beetle, town

longhorn beetle

view more common names online...

EPPO Categorization: A2 list

view more categorizations online...

EPPO Code: AELSSA


CT20H C

@ 2020-05-26

GEOGRAPHICAL DISTRIBUTION

@ 2020-05-26

The area of origin of *T. sartus* is thought to be Pakistan and Western India, from which it spread westwards to Afghanistan and Iran and northwards to the Central Asian countries; Tajikistan, Kyrgyzstan, Turkmenistan, Uzbekistan where it was first found in 1911 (in Samarkand, UZ). The pest continues to increase its range in these countries (Orlinski *et al.*, 1991; EPPO, 2005). *T. sartus* was also found in several poplar trees in South Kazakhstan (Shymkent), however, the researcher believes that the pest will not be able to survive there (Kostin, 1973). There is some information about findings in Japan, Malaysia and Sri Lanka (CAPS Program Resource, 2020) but more data is needed to confirm these findings. *T. sartus* is found usually in lower mountainous areas up to an altitude of 2000 m.



Dynamic parts: automatically generated by GD



EPPO's achievements are only possible because of the collaboration of experts from our region but also from other parts of the world. Thanks to all!



All EPPO Standards and recommendations are available in Global Database

