



Food and Agriculture
Organization of the
United Nations



International
Plant Protection
Convention

INTERNATIONAL STANDARD FOR PHYTOSANITARY MEASURES 28

PHYTOSANITARY TREATMENT

ISPM 28
ANNEX 33

ENG

PT 33: Irradiation treatment for *Bactrocera dorsalis*

Produced by the Secretariat of the
International Plant Protection Convention (IPPC)

This page is intentionally left blank

This phytosanitary treatment was adopted by the Fifteenth Session of the Commission on Phytosanitary Measures in 2021.
The annex is a prescriptive part of ISPM 28.

ISPM 28

Phytosanitary treatments for regulated pests

PT 33: Irradiation treatment for *Bactrocera dorsalis*

Adopted 2021; published 2021

Scope of the treatment

This treatment describes the irradiation of fruits and vegetables at 116 Gy minimum absorbed dose to prevent the emergence of adults of *Bactrocera dorsalis* at the stated efficacy.¹

Treatment description

Name of treatment	Irradiation treatment for <i>Bactrocera dorsalis</i>
Active ingredient	n/a
Treatment type	Irradiation
Target pest	<i>Bactrocera dorsalis</i> (Hendel, 1912) (Diptera: Tephritidae)
Target regulated articles	All fruits and vegetables that are hosts of <i>Bactrocera dorsalis</i>

Treatment schedule

Minimum absorbed dose of 116 Gy to prevent the emergence of adults of *Bactrocera dorsalis*.

There is 95% confidence that the treatment according to this schedule prevents emergence of the adult stage from not less than 99.9963% of eggs and larvae of *Bactrocera dorsalis*.

This treatment should be applied in accordance with the requirements of ISPM 18 (*Guidelines for the use of irradiation as a phytosanitary measure*).

Other relevant information

Because irradiation may not result in outright mortality, inspectors may encounter live but non-viable *Bactrocera dorsalis* (eggs, larvae or puparia) during the inspection process. This does not imply a failure of the treatment.

The Technical Panel on Phytosanitary Treatments based its evaluation of this treatment on the research reported by Zhao *et al.* (2017), which determined the efficacy of irradiation as a treatment for this pest in *Psidium guajava*. In addition, the work of Follett and Armstrong (2004) supports this schedule.

The efficacy of this schedule was calculated based on a total of 100 684 third-instar larvae treated with no adult emergence; the control emergence was 81%.

¹ The scope of phytosanitary treatments does not include issues related to pesticide registration or other domestic requirements for contracting parties' approval of treatments. Treatments adopted by the Commission on Phytosanitary Measures may not provide information on specific effects on human health or food safety, which should be addressed using domestic procedures before contracting parties approve a treatment. In addition, potential effects of treatments on product quality are considered for some host commodities before their international adoption. However, evaluation of any effects of a treatment on the quality of commodities may require additional consideration. There is no obligation for a contracting party to approve, register or adopt the treatments for use in its territory.

Extrapolation of treatment efficacy to all fruits and vegetables was based on knowledge and experience that radiation dosimetry systems measure the actual radiation dose absorbed by the target pest independent of host commodity, and evidence from research studies on a variety of pests and commodities. These include studies on the following pests and hosts: *Anastrepha fraterculus* (*Eugenia pyriformis*, *Malus pumila* and *Mangifera indica*), *Anastrepha ludens* (*Citrus paradisi*, *Citrus sinensis*, *Mangifera indica* and artificial diet), *Anastrepha obliqua* (*Averrhoa carambola*, *C. sinensis* and *Psidium guajava*), *Anastrepha suspensa* (*Averrhoa carambola*, *C. paradisi* and *Mangifera indica*), *Bactrocera tryoni* (*C. sinensis*, *Solanum lycopersicum*, *Malus pumila*, *Mangifera indica*, *Persea americana* and *Prunus avium*), *Cydia pomonella* (*Malus pumila* and artificial diet), *Grapholita molesta* (*Malus pumila* and artificial diet), *Pseudococcus jackbeardsleyi* (*Cucurbita* sp. and *Solanum tuberosum*) and *Tribolium confusum* (*Triticum aestivum*, *Hordeum vulgare* and *Zea mays*) (Bustos *et al.*, 2004; Gould and von Windeguth, 1991; Hallman, 2004a, 2004b, 2013; Hallman and Martinez, 2001; Hallman *et al.*, 2010; Jessup *et al.*, 1992; Mansour, 2003; Tunçbilek and Kansu, 1996; von Windeguth, 1986; von Windeguth and Ismail, 1987; Zhan *et al.*, 2016). It is recognized, however, that treatment efficacy has not been tested for all potential fruit and vegetable hosts of the target pest. If evidence becomes available to show that the extrapolation of the treatment to cover all hosts of this pest is incorrect, the treatment will be reviewed.

References

The present annex may refer to ISPMs. ISPMs are available on the International Phytosanitary Portal (IPP) at <https://www.ippc.int/core-activities/standards-setting/ispms>.

- Bustos, M.E., Enkerlin, W., Reyes, J. & Toledo, J.** 2004. Irradiation of mangoes as a postharvest quarantine treatment for fruit flies (Diptera: Tephritidae). *Journal of Economic Entomology*, 97: 286–292.
- Follett, P.A. & Armstrong, J.W.** 2004. Revised irradiation doses to control melon fly, Mediterranean fruit fly, and oriental fruit fly (Diptera: Tephritidae) and a generic dose for tephritid fruit flies. *Journal of Economic Entomology*, 97(4): 1254–1262.
- Gould, W.P. & von Windeguth, D.L.** 1991. Gamma irradiation as a quarantine treatment for carambolas infested with Caribbean fruit flies. *Florida Entomologist*, 74: 297–300.
- Hallman, G.J.** 2004a. Ionizing irradiation quarantine treatment against oriental fruit moth (Lepidoptera: Tortricidae) in ambient and hypoxic atmospheres. *Journal of Economic Entomology*, 97: 824–827.
- Hallman, G.J.** 2004b. Irradiation disinfestation of apple maggot (Diptera: Tephritidae) in hypoxic and low-temperature storage. *Journal of Economic Entomology*, 97: 1245–1248.
- Hallman G.J.** 2013. Rationale for a generic phytosanitary irradiation dose of 70 Gy for the genus *Antastrepha* (Diptera: Tephritidae). *Florida Entomologist*, 96(3): 983–990.
- Hallman, G.J., Levang-Brilz, N.M., Zettler, J.L. & Winborne, I.C.** 2010. Factors affecting ionizing radiation phytosanitary treatments, and implications for research and generic treatments. *Journal of Economic Entomology*, 103: 1950–1963.
- Hallman, G.J. & Martinez, L.R.** 2001. Ionizing irradiation quarantine treatment against Mexican fruit fly (Diptera: Tephritidae) in citrus fruits. *Postharvest Biology and Technology*, 23: 71–77.
- Jessup, A.J., Rigney, C.J., Millar, A., Sloggett, R.F. & Quinn, N.M.** 1992. Gamma irradiation as a commodity treatment against the Queensland fruit fly in fresh fruit. In: *Use of irradiation as a quarantine treatment of food and agricultural commodities*. Proceedings of the Final Research Coordination Meeting on Use of Irradiation as a Quarantine Treatment of Food and Agricultural Commodities, Kuala Lumpur, August 1990, pp. 13–42. Vienna, International Atomic Energy Agency.
- Mansour, M.** 2003. Gamma irradiation as a quarantine treatment for apples infested by codling moth (Lepidoptera: Tortricidae). *Journal of Applied Entomology*, 127: 137–141.

- Tunçbilek, A.S. & Kansu, I.A.** 1996. The influence of rearing medium on the irradiation sensitivity of eggs and larvae of the flour beetle, *Tribolium confusum* J. du Val. *Journal of Stored Products Research*, 32: 1–6.
- von Windeguth, D.L.** 1986. Gamma irradiation as a quarantine treatment for Caribbean fruit fly infested mangos. *Proceedings of the Florida State Horticultural Society*, 99: 131–134.
- von Windeguth, D.L. & Ismail, M.A.** 1987. Gamma irradiation as a quarantine treatment for Florida grapefruit infested with Caribbean fruit fly, *Anastrepha suspensa* (Loew). *Proceedings of the Florida State Horticultural Society*, 100: 5–7.
- Zhan, G., Shao, Y., Yu, Q., Xu, L., Liu, B., Wang, Y. & Wang, Q.** 2016. Phytosanitary irradiation of Jack Beardsley mealybug (Hemiptera: Pseudococcidae) females on rambutan (Sapindales: Sapindaceae) fruits. *Florida Entomologist*, 99 (Special Issue 2): 114–120.
- Zhao, J., Ma, J., Wu, M., Jiao, X., Wang, Z., Liang, F. & Zhan, G.** 2017. Gamma radiation as a phytosanitary treatment against larvae and pupae of *Bactrocera dorsalis* (Diptera: Tephritidae) in guava fruits. *Food Control*, 72: 360–366.

Publication history

This is not an official part of the standard

- 2017-06 Treatment submitted in response to 2017-02 call for treatments.
- 2018-01 Technical Panel on Phytosanitary Treatments (TPPT) reviewed and requested further information from submitter.
- 2018-04 Submitter supplied additional information.
- 2018-05 Standards Committee (SC) added the topic *Irradiation treatment for oriental fruit fly Bactrocera dorsalis on all fresh commodities* (2017-015) to the TPPT work programme.
- 2018-06 TPPT revised the draft and recommended to SC for consultation.
- 2018-11 TPPT final review via e-forum (2018_eTPPT_Oct_02).
- 2019-01 SC approved the draft for consultation via e-decision (2019_eSC_May_06).
- 2019-07 First consultation.
- 2020-02 TPPT reviewed and approved the responses to consultation comments and recommended the draft for second consultation.
- 2020-06 SC approved for second consultation via e-decision (2020_eSC_May_22).
- 2020-07 Second consultation.
- 2020-11 TPPT virtual meeting recommended to SC for approval for adoption by the CPM.
- 2021-03 CPM-15 adopted the phytosanitary treatment.
- ISPM 28. Annex 33.** *Irradiation treatment for Bactrocera dorsalis* (2021). Rome, IPPC, FAO.
- 2021-04 IPPC Secretariat applied ink amendments as noted by CPM-15 (2021).

Publication history last updated: 2021-04

IPPC

The International Plant Protection Convention (IPPC) is an international plant health agreement that aims to protect global plant resources and facilitate safe trade.

The IPPC vision is that all countries have the capacity to implement harmonized measures to prevent pest introductions and spread, and minimize the impacts of pests on food security, trade, economic growth, and the environment.

Organization

- ◆ There are over 180 IPPC contracting parties.
- ◆ Each contracting party has a national plant protection organization (NPPO) and an Official IPPC contact point.
- ◆ 10 regional plant protection organizations (RPPOs) have been established to coordinate NPPOs in various regions of the world.
- ◆ IPPC liaises with relevant international organizations to help build regional and national capacities.
- ◆ The Secretariat is provided by the Food and Agriculture Organization of the United Nations (FAO).

International Plant Protection Convention Secretariat
ippc@fao.org | www.ippc.int

Food and Agriculture Organization of the United Nations
Rome, Italy

